

Geophysical Trends inferred from 20 years of AIRS infrared global observations

S. DeSouza-Machado¹, L. Larrabee Strow^{1,2}, R. Kramer³

¹JCET/GESTAR2, University of Maryland Baltimore County, Baltimore, Maryland

²Dept of Physics, University of Maryland Baltimore County, Baltimore, Maryland

³NOAA GFDL, Princeton New Jersey

Key Points:

- The 20+ year radiance record of NASA's AIRS sounder contains detailed vertical information about changes in geophysical parameters.
- ~~A novel method retrieves~~ We use an algorithm uniquely designed to retrieve geophysical trends from the radiance observation record, using stable channels ~~in radiance space and straightforward and traceable a-priori~~.
- Comparisons are made to trends from monthly ~~Numerical Weather Prediction~~ re-analysis fields and L3 operational data products.

Corresponding author: Sergio DeSouza-Machado, sergio@umbc.edu

Abstract

Daily spectral radiance observations by NASA’s Atmospheric Infrared Sounder contain detailed information about surface and atmospheric temperature and water vapor. We obtain climate geophysical trends from 20 years (2002/09-2022/08) of AIRS observations using a novel method operating mostly in radiance space. The observations are binned into 3×5 degree tiles using 16 day intervals, after which nominally clear scenes are selected for each tile to construct the spectral radiance time series. De-seasonalized spectral trends are then obtained, which are inverted using a physical retrieval to obtain geophysical trends. This approach is distinct from traditional use of radiances whereby trends are generated after operational retrievals or assimilation into Numerical Weather Prediction models. Our approach rigorously ties the derived geophysical trends to the observed radiance trends, using far fewer computational resources and time. The retrieved trends are compared to trends derived from ERA5 and MERRA2 reanalysis model fields, and NASA Level3 AIRS v7 and CLIMCAPS v2 data. Our retrieved surface temperature trends agree quite well with ERA5, CLIMCAPS and the GISS surface climatology trends. Atmospheric temperature profile trends exhibit some variability amongst all these data sets, especially in the polar stratosphere. Water vapor profile trends are nominally similar among the data sets except for the AIRS v7 which exhibits drying trends in the mid troposphere. Spectral closure between observed trends and those computed by running the reanalysis and NASA L3 monthly fields through a radiative transfer code are discussed, with the major differences arising in the water vapor sounding region.

Plain Language Summary

The current generation of infrared sounders, designed for weather forecasting purposes, have been ~~in orbit around the Earth~~ operational for a long enough time to enable anomaly and trending studies for climate purposes. ~~Traditionally their daily obtained radiance data has been~~ The daily radiance observations are routinely used for operational atmospheric state retrievals ~~, or assimilation into Numerical Weather Prediction and assimilation into reanalysis~~ models, after which climate anomaly studies are ~~made~~. ~~In this paper we use the raw radiance spectral data to form radiances~~ enabled. ~~Here we use a purpose built algorithm to directly turn radiance observations into geophysical~~ anomalies and trends ~~, after which we do a one step atmospheric state retrieval. This novel approach has the benefit of using only stable channels together with~~ with full error characterization. This unique approach for observational climate trending uses only stable low noise sounding channels, easily understood assumptions and well tested retrieval algorithms ~~to do the trend or anomaly geophysical retrieval, which has full error characterization.~~

1 Introduction

1 Introduction

On time scales of a few hours to a week, accurate time resolved, global measurements of atmospheric water vapor amounts and temperatures are necessary for short to medium range weather forecasting. Non condensing Greenhouse Gases (GHG) such as CO_2 live in the atmosphere for relatively long times (on the order of 10-100 years). At these longer timescales, the gradually increasing opacity of the atmosphere (in the $500 - 800 \text{ cm}^{-1}$ ($15\text{-}20 \mu\text{m}$) infrared region of the electromagnetic spectrum) due to increasing CO_2 amounts ($\sim 2.2 \text{ ppmv/year}$ (Keeling et al., 1976)) and other long lived GHG serves as a controlling knob mechanism (Muller et al., 2016) that can slowly increase the surface and air temperatures. A timeline summary of pioneering scientific work recognizing the contribution of GHG to Earth temperatures is found in (Anderson et al., 2016), including Joseph Fourier (1827) understanding the opacity of atmosphere to infrared radiation, John Tyndall (1872) recognizing infrared absorption by H_2O and CO_2 , a prediction of rising surface temperatures from increasing GHG concentrations (Arrhenius,

1896), to Guy Callendar (1958) correlating land surface temperature changes with increases in CO₂. A key paper (Manabe & Wetherald, 1967) estimated a 2 K temperature increase of an atmosphere (with fixed relative humidity) for a CO₂ doubling. An example of a more recent relevant paper modeling how CO₂ increases change the atmospheric opacity is Jevanjee et al. (2021). Note the changing concentrations of other non condensing GHG such as N₂O and CH₄ together with CO₂ also change the opacities in other regions of the thermal and shortwave infrared, but they can be of diminished importance due to the nonlinear Planck function which peaks at about 15 μm for the 250-300 K temperatures typical of the Earth atmosphere.

The higher temperatures from these and other external forcings increase the amount of water vapor that the atmosphere can hold. Water vapor itself is a GHG (Held & Soden, 2000; Muller et al., 2016) and serves to further increase the opacity of the atmosphere, both in the Far Infrared (0-500 cm⁻¹) and in the thermal infrared window region (800-1200 cm⁻¹) and the water vapor sounding region (1350 - 2000 cm⁻¹). Atmospheric water vapor amounts are highly variable in time and space. Though the lifetime of water in the atmosphere (the hydrological cycle of evaporation from surface, condensation into a cloud, precipitation back to the surface) is on the order of less than two weeks (van der Ent & Tuinenburg, 2017; Sodemann, 2019), the additional radiative forcing is a positive feedback which amplifies the temperature changes due to the long-lived GHG (Manabe & Wetherald, 1967; Muller et al., 2016). This also has implications for precipitation changes (Held & Soden, 2006).

There is therefore a need to have an accurate, global, high vertical resolution long term time series of temperature and water vapor measurements since they will provide a critical record of climate change, which will help scientists understand climate change by providing evidence based comparisons to outputs of climate models, as well as validate climate model predictions. Key documents which underscore the importance of accurate measurements of global temperature and humidity include the Intergovernmental Panel on Climate Change (IPCC) reports; see for example (Houghton et al., 1990; IPCC, 2021).

In this paper we focus on surface temperature, and atmospheric temperature and water vapor trends. Climate monitoring traditionally uses averages computed over 30 consecutive years together with quantifying the linear trends (Thorne et al., 2011; Scherrer et al., 2024). Monitoring observed short term trends (~ 20 -30 years) is possibly preferable to longer term trends (~ 50 -100 years) due to nonlinearities in the physical models (Scherrer et al., 2024), and can help identify which regions are warming faster than others, confirm whether estimates from models agree with the observations, and identify physical processes or regions where models need to be improved (for example cloud processes, natural variability due to volcanic eruptions and natural oscillations in the Earth system). Temperature increases affect the environment including for example sea level rises and melting of glaciers, and impact society by for example affecting crop yields and inducing heat related illnesses. Climate models predict a fingerprint of vertical trend profiles together with uncertainties (Thorne et al., 2011) which can be compared against trends and uncertainties from observational data. Water vapor trends predicted by models can similarly be fingerprinted and compared against observations (Allan et al., 2022). Changes in atmospheric water vapor leads to radiative forcings (Dessler et al., 2008), changes in precipitation amounts, and through the release of latent heat as the vapor condenses into clouds, affect the distribution of heat and planetary circulation (Schneider & Levine, 2010).

Observational data comes from two main sources. The first is ground based instruments, a prime example of which are radiosondes (Durre et al., 2006), which typically record temperatures, windspeeds and humidity measurements as a function of pressure. The data is available from the 1960s, but while these instruments can be regularly calibrated, their coverage is sparse, typically over land and very little over ocean (though there is an extensive network of buoys to measure ocean temperature and salinity pro-

files (Wong et al., 2020)). On the other hand satellite based observations covering the Earth started in the 1970s and are now available across almost the entire electromagnetic spectrum, from passive microwave and infrared sounders that provide global information about temperature and humidity 24 hours a day, to passive visible and ultraviolet imagers that provide global information about clouds and aerosols during daytime conditions, to active (lidar and radar) instruments that provide cloud and aerosol profile information with more limited coverage, to radio occultation instruments which use radio signals from the Global Positioning Satellites (GPS) to obtain temperature and humidity information. Each of these satellite instruments provide valuable information about the atmosphere. In this paper we use 20 years of observational data (September 2002 - August 2022) from the Atmospheric Infrared Sounder (AIRS), which is a new generation infrared nadir sounder.

Model data can come from re-analysis (Kalnay et al., 1996; Gelaro & Coauthors, 2017; Hersbach et al., 2020) or climate models (Eyring et al., 2016). In this paper we use re-analysis model data, which ingest observations from a wide variety of instruments. Depending on the re-analysis model, the assimilation+dynamical system may give inaccurate trends because of one or more of the following reasons (Cai & Kalnay, 2004; Kistler et al., 2001; Shao et al., 2023) : they typically use observing systems that change with time as new *in-situ* or satellite based instruments become operational, the observations may have biases and errors that are not correctly accounted for, the assimilated observations may be obtained (or simply not available) in cloud conditions, the dynamical models cannot contain all the physics (eg land topography, cloud microphysics) at the finite sized grid boxes used, and often do not use eg time varying concentrations of CO₂ or stratospheric aerosols. Modern re-analysis systems have mostly addressed these issues (Dee et al., 2011) but there still are concerns in for example data sparse regions (Hobbs et al., 2020; Bromwich et al., 2024). We note that the 20+ years of AIRS observations (2002-2022) used in this paper meant we chose to exclude comparing climate model data from the latest (sixth) Coupled Model Intercomparison Project (CMIP6) since it covers the period 1850-2014 (Eyring et al., 2016).

Another source of data used in this paper, also tied to observations, come from L2 or L3 retrievals. Operational NASA AIRS daily Level 2 products and monthly Level 3 products (derived from Level 2) used in this paper retrieve the atmospheric state using cloud-cleared radiances derived from a 3x3 grid of individual scenes. A main characteristic of traditional L2 retrievals is the requirement for a good *a-priori* state for each inversion, making errors in the *a-priori* difficult to distinguish from true variability in the observations. Measurements by visible imagers which have ~ 1 km horizontal resolution or better (King et al., 2013) suggest global cloud free fractions of $\sim 30\%$, but the 15 km footprint of typical sounders means at most 5% of the hyperspectral observations can be considered “cloud-free.” Current operational NASA L2 products for AIRS use the method of cloud clearing on observed radiances in partly cloudy scene conditions before doing the geophysical retrieval. The cloud clearing method takes in the raw observed allsky radiances and solves for an estimate of clear column radiances by examining adjacent Fields of View (FOVs) to estimate the cloud effects on the observations. The method assumes any differences are solely due to different cloud amounts in each FOV, and significantly increases geophysical retrieval yields (to about 50-60%) (Smith & Barnet, 2023). The resulting cloud cleared radiances (CCR), distinct from clear sky radiances which are obtained under nominally clear conditions, have increased noise especially in the lower atmosphere sounding channels; in addition the subsequent retrieval depends on the first guess (which is a neural net for AIRS v7 and MERRA2 reanalysis for CLIMCAPS v2). The reader is referred to AIRS L2 literature (Susskind et al., 2003; Susskind, 2006; Susskind et al., 2014; Smith & Barnet, 2020, 2023) for more details about cloud clearing and the L2 algorithms.

1.1 Using hyperspectral infrared radiance observations directly for climate trending

The “observational data \rightarrow sophisticated assimilation or retrieval algorithms \rightarrow daily or monthly products \rightarrow climate trends” approaches outlined in the previous paragraph are not tailored for climate anomalies or trending. Both reanalysis and Level 2 products require large computational resources, that preclude full dataset re-processing to help fully understand trends. In this paper we use an approach purpose-designed to produce climate anomalies and trends directly from infrared radiance observations. We work exclusively in radiance space and form either anomalies or trends from the underlying well characterized and understood radiances (Strow & DeSouza-Machado, 2020), in order to do a geophysical trend or anomaly retrieval. Moreover, our novel approach has zero temperature *a-priori* and minimal water vapor *a-priori*. This completely sidesteps time variability and the accuracy of the *a-priori* which causes errors in the retrievals, and ensures our work examines trends directly inferred from the radiances versus those from traditional methods. This leads to more unbiased results that directly highlight the conditions (for example stratospheric water vapor) where the sensor has limited sensitivity. The work presented here, once the averaged/sorted observations are available, can be processed in hours to days, and can be duplicated by small research groups with ease.

The Outgoing Longwave Radiation (OLR) balanced against the Incoming Solar Radiation at the top of the Earth’s atmosphere, is the fundamental driver of the climate system (Brindley & Bantges, 2016). Broadband measurements (integrated across the entire longwave, or entire shortwave bands) have been available for over 40 years, and provide a valuable record. However the single integrated measurement effectively smears out competing effects such as increases in OLR due to increases in surface temperatures versus decreases in OLR due to changing $\text{CO}_2, \text{H}_2\text{O}, \text{O}_3, \text{CH}_4$ Greenhouse Gas (GHG) concentrations (Brindley & Bantges, 2016). Passive microwave and infrared instruments with handfuls of channels have also been flown since the late 1970s for meteorological purposes. For example the Microwave Sounding Unit (MSU) and Advanced Microwave Sounding Unit (AMSU) provide global scale records of upper atmospheric temperatures (Mears & Wentz, 2009, 2016). Another example is the 20 channel High resolution Infrared Radiation Sounder (see for example (Harries et al., 1998; Shi & Bates, 2011; Menzel et al., 2016)), which provides an $\sim 40+$ year global observational dataset. The advantage of these instruments is their spectrally resolved channels are capable of providing radiance measurements from which one can quantify the effects of individual GHG and surface/air temperature changes. Limitations with these observational records include drifts of the orbits or instruments, inter-calibrating the individual instruments contributing to the record (which individually have lifetimes of the order of 5-10 years), and the spectrally wide channels mean the vertical weighting functions are very broad which only allows for limited vertical resolution (typically a few kilometers); the same wide channels also typically include the radiative effects of more than one gas. All these serve to constrain climate studies. For example the upper tropospheric water vapor sensitivity of the HIRS instruments are typically deep layers between 200-500 mb (see Muller et al. (2016) and references therein), while the water vapor feedback is very sensitive to the H_2O changes in the thinner layers extending between the cold dry upper troposphere and lower stratosphere (Muller et al., 2016).

These limitations have largely been mitigated by the new generation of infrared sounders, which have high spectral resolution (superior vertical resolution), are very stable and whose overlapping orbits and long lifetimes allows for continual inter-calibration and monitoring of the stability of these instruments; see for example (Strow et al., 2021). The first of the new generation of low noise, high stability hyperspectral sounders is NASA’s Atmospheric Infrared Sounder (AIRS). The instrument has been in continuous operation since September 2002, making Top of Atmosphere (TOA) radiance observations at a typical 15km (at nadir) horizontal resolution. Follow on instruments with similar charac-

teristics and abilities include the ESA’s Infrared Atmospheric Sounding Interferometer (IASI) and NOAA’s Cross-track Infrared Sounder (CrIS), operational since June 2007 and March 2012 respectively. The latter two already have follow on missions planned till the 2040s, and together these three sounders will provide scientists with a 40 year high quality, near continuous observational dataset for climate studies.

Infrared radiances contain a wealth of information. A short list focusing on scientific contributions using the AIRS radiance observations includes improvements in surface temperature, atmospheric temperature and water vapor profiles in weather forecasting models (see for example LeMarshall et al. (2006); Andersson et al. (2007)), retrieving mixing ratios of greenhouse gases such as CO₂ (Chedin et al., 2005), CH₄ (Zou et al., 2019) and O₃ (Fu et al., 2018). Clouds (Kahn et al., 2005, 2014) and large aerosols (volcanic ash and dust) (Carn et al., 2005; De Souza-Machado et al., 2010) can also be detected and quantified. Examples of other trace gases that can be detected and quantified are CO (McMillan et al., 2005) and NH₃ (Warner et al., 2016). This list is not exhaustive and in addition multiple papers have similarly been published using CrIS and IASI data.

The stability and accuracy of the AIRS instrument is documented in recent work on analyzing 16 years of AIRS radiance anomalies over cloud-free ocean (Strow & DeSouza-Machado, 2020). Geophysical retrievals on the anomalies yielded CO₂, CH₄, N₂O and surface temperature time series that compared well against in-situ NOAA Global Monitoring Laboratories (GML) trace gas measurements and NOAA Goddard Institute of Space Studies (GISS) surface temperature data respectively. A significant difference between this paper and (Strow & DeSouza-Machado, 2020) is the nominally clear scenes used in this paper are selected uniformly from all over the Earth, while the clear scenes in the latter were zonal averages which were sometimes concentrated in certain regions. Here we derive geophysical trends from 20 years (September 2002 - August 2022) of AIRS measurements over $\sim 3 \times 5$ degree tiles covering the Earth, chosen such that the number of observations in each tile is roughly equal. A companion paper will utilize the geophysical trend results to derive Outgoing Longwave Radiation (OLR) trends and non-local clearsky feedback parameters. Nominally clear scenes for each tile are picked out using a quantile approach; from the time series, radiances trends are made over the entire Earth, from which geophysical trends are retrieved.

Observed infrared spectral trends from AIRS has already been a focus of earlier work by (X. Huang et al., 2023) who studied a slightly shorter time period (2002-2020) using the nadir L1B radiance observations (which have no or minimal frequency corrections compared to the L1C radiance dataset we use here). Similarly the paper by (Raghuraman et al., 2023) converted the AIRS observed radiances to Outgoing Longwave radiation (OLR) in the 0-2000 cm⁻¹ range, but neither of these studies involve retrieving geophysical trends from radiance spectral trends. Instead they include the effects of GHG forcings and convert various model trends (such as ERA5) to spectral trends for comparison against the observed spectral trends, which we also show in Appendix B. Teixeira et al. (2024) used the AIRS observations between 2003-2012 to measure the impact of increased CO₂ on the outgoing longwave radiation. Another noteworthy examination of the time evolution of high spectral resolution infrared radiances (converted to spectral outgoing longwave radiation (OLR) fluxes) by Whitburn et al. (2021) covered 10 years (2007-2017) of IASI observations. They confirmed that the IASI-derived fluxes agreed well with increases in GHG gas concentrations and El-Nino Southern Oscillation (ENSO) events within that time frame. A more recent paper (Roemer et al., 2023) used the 10 year IASI observations to derive allsky longwave feedback spectral components (water vapor, CO₂, window, ozone) and total values, while also estimating clear sky feedback values. Other relevant studies involving high spectral resolution infrared measurements include the allsky interannual variability at different spatial scales using 5 years (2007-

2012) of IASI observations (Brindley et al., 2015), and comparing Global Climate Model simulations to AIRS radiances as a diagnostic of model biases (Y. Huang et al., 2007).

We will refer to our results as the AIRS Radiance Trends (AIRS_RT). Comparisons are made against monthly output from the European Center for Medium Weather Forecast fifth generation reanalysis (ERA5) (Hersbach et al., 2020) and NASA’s second generation Modern-Era Retrospective analysis for Research and Applications (MERRA2) (Gelaro & Coauthors, 2017), and also against the official monthly AIRS L3 products which are AIRS v7 L3 (Susskind et al., 2014; Tian et al., 2020) and CLIMCAPS v2 L3 (Smith & Barnet, 2019, 2020). Detailed geophysical trends and spectral closure studies are presented for the averaged ascending (daytime (D)) and descending (nighttime (N)) trends; the appendix briefly discusses separate D and N trends.

~~NASA’s Atmospheric Infrared Sounder (AIRS) became operational in September 2002, as the first of the new generation of low noise, high stability hyperspectral sounders, making Top of Atmosphere (TOA) radiance observations at a typical 15km (at nadir) horizontal resolution. Follow on instruments with similar characteristics and abilities include the ESA’s Infrared Atmospheric Sounding Interferometer (IASI) and NOAA’s Cross Track Infrared Sounder (CrIS), operational since June 2007 and March 2012 respectively. The latter two already have follow on missions planned till the 2040s, and together these three sounders will provide scientists with a 40 year high quality, near continuous observational dataset for climate anomaly and trending studies.~~

~~Infrared radiances contain a wealth of information, including but not limited to surface temperature, atmospheric temperature and water amount, and mixing ratios of greenhouse gases such as carbon dioxide, and . Measurements by visible imagers which have ~ 1 km horizontal resolution or better suggest global cloud free fractions of $\sim 30\%$, but the 15 km footprint of typical sounders means at most 5% of the hyperspectral observations can be considered “cloud-free.” Current operational NASA L2 products use the method of cloud clearing on observed radiances in partly cloudy scene conditions before doing the geophysical retrieval. The cloud clearing method solves for an estimate of clear column radiances by examining adjacent Fields of View (FOVs) to estimate the cloud effects on observed allsky radiances, assuming any differences are solely due to different cloud amounts in each FOV, and significantly increases geophysical retrieval yields (to about 50-60%) . This does introduce increased noise in the cloud cleared radiances of the lower atmosphere sounding channels; in addition the subsequent retrieval depends on the first guess (which is a neural net for AIRS v7 and MERRA2 reanalysis for CLIMCAPS v2). The reader is referred to for more details.~~

~~In this paper we work directly in radiance space and form either anomalies or trends from the underlying well characterized and understood radiances, in order to do a geophysical trend or anomaly retrieval. The work presented here, once the averaged/sorted data is available, can be processed in hours to days, and can be duplicated by small research groups with ease. Moreover, our novel approach has zero temperature *a-priori* and minimal water vapor *a-priori*. This completely sidesteps time variability and the accuracy of the *a-priori* which causes errors in the retrievals, and ensures our work examines trends directly inferred from the radiances versus those from traditional methods. This leads to more unbiased results that directly highlight the conditions (for example stratospheric water vapor) where the sensor has limited sensitivity.~~

~~The approaches used in this work are therefore very different than climate anomalies or trends from reanalysis products or traditional Level 2 retrievals, neither of which are tailored for climate trends. Reanalysis products assimilate individual sensor scenes from many different instruments, and may have discontinuities as different instruments come online or go offline. Traditional Level 2 (and Level 3 products derived from Level 2) retrieve the atmospheric state for individual scenes (or effective cloud-cleared radiance derived from a 3x3 grid of individual scenes). Both reanalysis and Level 2 products require large~~

computational resources, that preclude full-dataset re-processing to help fully understand trends. A main characteristic of traditional L2 retrievals is the requirement for a good state for each inversion, making errors in the difficult to distinguish from true variability in the data, especially with regard to trends.

The stability and accuracy of the AIRS instrument is documented in recent work on analyzing 16 years of AIRS radiance anomalies over cloud-free ocean. Geophysical retrievals on the anomalies yielded K , $^{\circ}\text{C}$, and surface temperature time series that compared well against in-situ data from NOAA Global Monitoring Laboratories (GML) trace gas measurements and NOAA Goddard Institute of Space Studies (GISS) surface temperature data respectively. A significant difference between this paper and is the nominally clear scenes used in this paper are selected uniformly from all over the Earth, while the clear scenes in the latter were zonal averages which were sometimes concentrated in certain regions.

In this paper we expand upon our initial zonal clear sky analysis, to derive geophysical trends from 20 years (September 2002 – August 2022) of AIRS measurements over $\sim 3^{\circ} \times 5^{\circ}$ degree tiles covering the Earth, chosen such that the number of observations in each tile is roughly equal. An important concept introduced is spectral closure, whereby the observed clear sky spectral radiance trends are compared to spectral trends produced by running the monthly reanalysis or official NASA retrieved AIRS L3 products through an accurate clear sky radiative transfer code; close agreement in different sounding regions (such as 640-800 for temperature and $^{\circ}\text{C}$, 1350-1640 for water vapor, 1000-1150 for $^{\circ}\text{C}$) between the computed and actual observed spectral trends imply that trends from those geophysical parameters used in the computations are realistic while disagreement suggests otherwise. A companion paper will utilize the geophysical trend results to derive Outgoing Longwave Radiation (OLR) trends and non-local clearsky feedback parameters. Nominally clear scenes for each tile are picked out using a quantile approach; from the time series, radiances trends are made over the entire Earth, from which geophysical trends are retrieved.

Observed infrared spectral trends from AIRS has already been a focus of earlier work by who studied a slightly shorter time period (2002-2020) while converted the radiances to Outgoing Longwave radiation (OLR), but neither study involve retrievals from spectral trends to geophysical trends. Instead they convert various model trends (such as ERA5) to spectral trends and compare against the observed spectral trends. Our earlier work shows we can accurately account for the effects of GHG forcings Strow et al. (2021). In this paper we remove these GHG forcings from the observed AIRS spectral trends to concentrate on atmospheric temperature and water vapor and surface temperature, while the papers by include the GHG forcings in the model generated spectral trends. Another noteworthy examination of the time evolution of high spectral resolution infrared radiances (converted to spectral outgoing longwave radiation (OLR) fluxes) by Whitburn et al. (2021) covered 10 years (2007-2017) of IASI observations. They confirmed that the IASI-derived fluxes agreed well with increases in GHG gas concentrations and El-Nino Southern Oscillation (ENSO) events within that time frame. A more recent paper used the 10 year IASI data to derive allsky longwave feedback spectral components (water vapor, $^{\circ}\text{C}$, window, ozone) and total values, while also estimating clear sky feedback values.

We will refer to our results as the AIRS Radiance Trends (K). Comparisons are made against monthly output from the European Center for Medium Weather Forecast fifth generation reanalysis (ERA5) and NASA's second generation Modern-Era Retrospective analysis for Research and Applications (MERRA2), and also against the official monthly AIRS L3 products which are AIRS v7 L3 and CLIMCAPS v2 L3. Detailed geophysical trends and spectral closure studies are presented for the averaged ascending (daytime (D)) and descending (nighttime (N)) trends; the appendix briefly discusses separate D and N trends.

2 Datasets used in this study

Three main types of datasets are used in this study. The first is the AIRS L1C radiance ~~dataset we analyzed~~ observation dataset we analyze for this paper, which has both daytime (D) and nighttime (N) (ascending and descending) views of the planet. Second is the monthly operational L3 retrieval data, which are the AIRS v7 and the CLIMCAPS v2 products, also separated into D/N ~~data~~ subsets. Finally we also compared to trends from ERA5 and MERRA2 monthly reanalysis model fields. The ERA5 monthly dataset is available in 8 averaged time steps, so we match to the average AIRS overpass times and ~~compute~~ separate into (D/N) ~~data~~ sets over the 20 years, while MERRA2 monthly model fields are only available as one time step; included here for completeness we mention the NASA GISS surface temperature dataset, which like MERRA2 is only available as a monthly mean. This means four of the datasets : AIRS_RT (from AIRS L1C), AIRS L3 and CLIMCAPS L3, and ERA5 are separable into D/N, while the other two (MERRA2 and GISS) are only available as a diurnal averaged value. We describe these datasets in more detail below.

2.1 The AIRS instrument and L1C observational dataset

The Atmospheric Infrared Sounder (AIRS) on board NASA’s polar orbiting EOS/Aqua platform has 2378 channels, covering the Thermal Infrared (TIR) spectral range (roughly $649\text{--}1613\text{ cm}^{-1}$) and shortwave infrared ($2181\text{--}2665\text{ cm}^{-1}$). The full widths at half maximum satisfy $\nu/\delta\nu \sim 1200$. The (spectral dependent) noise is typically $\leq 0.2\text{K}$. The original L1b radiance ~~dataset~~ observations suffers from spectral gaps and noise ~~contaminated~~ data-contamination as detectors slowly fail. These limitations are addressed using a 2645 L1c channel observational dataset, where spectral gaps and some of the noise “pops” are filled in using principal component reconstruction (Manning et al., 2020) and is the dataset used to subset radiances analyzed in this paper. However we note that the results described in this paper used only the actual observed radiances in pristine, stable channels described in (Strow et al., 2021) and none of the synthetic channels. The Aqua platform is a polar orbiting satellite with 1.30 am descending (night time over equator) and 1.30 pm ascending (daytime over equator) tracks. Each orbit takes about 90 minutes, with the 16 passes yielding almost twice daily coverage of the entire planet. About ~ 3 million AIRS spectral observations have been obtained daily since AIRS became operational in late August 2002. The instrument has provided ~~data~~ observations almost continuously since then though there have been some shutdowns (each spanning a few days) such as during solar flare events.

In this paper we use the re-calibrated 2645 channel L1C radiance ~~data~~ observations (Strow & DeSouza-Machado, 2020) instead of the 2378 L1B ~~data~~ radiance observations. 20 years (spanning September 1, 2002–August 31, 2022) of AIRS L1C radiance ~~data is~~ observations are gridded into 4608 tiles covering the Earth : 72 longitude boxes which are all 5° in width, and 64 latitude boxes which are approximately 2.5° in width at the tropics but wider at the poles to keep the number of observations per 16 day intervals (which is the repeat cycle of the AIRS orbit on the Aqua satellite) roughly the same. This way there are ~ 12000 observations per 16 days per tile, which are roughly equally divided between the ascending/daytime (D) and descending/nighttime (N) tracks. In this paper we discuss results for both the ascending and descending tracks using a retrieval based on the longwave (LW) and midwave (MW) regions of the spectrum ($640\text{--}1620\text{ cm}^{-1}$ or $6\text{--}15\text{ }\mu\text{m}$).

In this paper our trend retrievals use only the AIRS channels that are stable in time, as quantified in (Strow et al., 2021). For example the shortwave (SW) channels are drifting at a higher rate than the LW/MW channels, which can lead to incorrect surface temperature rates, and are avoided in this paper. Similarly there are many channels in the LW and MW whose detectors are drifting in time, and which are also not used here.

For example there are some higher wavenumber (shorter wavelength) channels past the ozone band which have a significant drift in time, possibly due to changes in the polarization of the scan mirror coating with time. Therefore compared to other AIRS operational products used in this paper, our results use channels that are demonstrated to have high stability (Strow et al., 2021). We do note that some of the observed drifts in the AIRS channels stabilized after 6 years, so their impact is reduced when looking at 20 year trends.

2.2 Reanalysis Model fields

The ERA5 fifth generation reanalysis product from the European Center for Medium Range Weather Forecasts is freely available on monthly timescales from the Copernicus Climate Data Store. This monthly dataset is output at 37 pressure levels at 0.25° horizontal resolution (Hersbach et al., 2020), which is further subdivided into eight 3-hour averages per month (corresponding to 00,03,06,...21 UTC). For each month from September 2002-August 2022 we downloaded the surface temperature and pressure fields, as well as atmospheric temperature, water vapor and ozone fields. These are then colocated to each tile center using 2D spatial interpolation, as well as time interpolated according to the average AIRS overpass time as a function of month. From the resulting monthly time-series of reanalysis model fields for each tile, we generated (a) thermodynamic trends for surface temperature, air temperature, water vapor and ozone model fields (b) a 20 year average thermodynamic profile in order to produce jacobians for the linear trend retrievals (c) by using the model fields as input to the clear sky SARTA radiative transfer code (Strow, Hannon, DeSouza-Machado, et al., 2003) a monthly time series of clear sky radiances for each tile was generated, from which we could compute radiance trends. ~~We did this-~~ The matching to ERA5 reanalysis was done for both the ascending and descending ~~datasets~~ observations.

The MERRA version 2 (MERRA2) re-analysis used in this paper is the second generation (Gelaro & Coauthors, 2017) product from NASA’s Global Modeling and Assimilation Office. The monthly data we use is available on 42 pressure levels at a horizontal resolution of $0.5^\circ \times 0.625^\circ$, but only one monthly mean diurnally averaged output is available per month. Similar to the ERA5 output, we colocated the MERRA2 surface temperature, atmospheric temperature, water vapor and ozone fields to our tile centers for each month starting September 2002 in order to produce a time series of radiance and model output, from which radiance and thermodynamic trends could be computed for comparisons against other datasets in this study; similar to above we also generated a monthly time series of clear sky radiances for each tile, from which we could compute clear sky radiance trends based on MERRA2.

The NASA Goddard Institute of Space Studies (GISS) ~~surface-temperature data~~ v4 surface temperature data (2023, 2005; Lenssen et al., 2019) is a monthly dataset based primarily on near surface temperatures land stations, and data from ships and buoys. As with MERRA2 we obtained one monthly mean dataset per month, which we could not separate into descending (N) or ascending (D) tracks.

2.3 AIRS L3 Products

NASA routinely produces two retrievals from the daily AIRS L1C ~~data-observed~~ each-day observations, which are AIRS v7 (Susskind et al., 2014; Tian et al., 2020) and CLIMCAPS v2 (Smith & Barnet, 2019, 2020). Both use the cloud clearing process but there are significant algorithmic differences; in particular the AIRS v7 product is initialized by a neural net, while CLIMCAPS uses MERRA2 for its initialization. The L2 products are then individually turned into L3 monthly products, for both the ascending (daytime) and descending (nighttime) observational data. The timeseries of thermodynamic profiles were used as input to the clear sky SARTA RTA to generate radiances, after which radiance trends and thermodynamic trends are also produced.

2.4 Other L3 Products

The Microwave Limb Sounder (MLS) monthly binned water vapor (H₂O) mixing ratio dataset (Livesey et al., 2006; Lambert et al., 2007, 2021), which contains ~~data at spatial coverage retrieved fields covering~~ $\pm 82^\circ$ latitude, at a spatial resolution of $4^\circ \times 5^\circ$ and useful vertical range between 316 and 0.00215 hPa was used in this paper to improve retrieval trends in the upper atmosphere.

3 Filtering the Observational Data for clear scenes

Here we discuss the “clear scene” selection from all the observed data stored for each of the 72×64 tiles. Ideally we would prefer to use a MODIS cloud fraction product (1 km) colocated to the 15 km AIRS footprints, but this is presently unavailable. Our earlier work used an uniform clear flag over ocean (Strow et al., 2021) which will not work well over land because of surface inhomogeneity. In this section we discuss an alternative clear filter based on the hottest 10 percent of AIRS observations that are present inside any 16 day tile, over any location.

3.1 Observed BT1231 Distributions

The radiances measured in thermal infrared window region ($800\text{--}1000\text{ cm}^{-1}$ and $1100\text{--}1250\text{ cm}^{-1}$) are dominated by the effects of the surface temperature, water vapor continuum absorption and cloud/aerosol effects. The effects of water vapor continuum absorption is largest in hot and humid tropical scenes (depressing the observations relative to surface temperatures by about 5-6 K, which reduces to about 2 K at $\pm 50^\circ$) and is almost negligible for cold, dry scenes (less than 1 K). Scattering and absorption by liquid and ice clouds also affects the window region (Deep Convective Clouds can depress the window channel observations by as much as 100 K relative to surface temperatures). For each tile, we use the 1231.3 cm^{-1} observation as our representative window channel (AIRS L1C channel ID = 1520), as it is minimally impacted by weak water vapor lines. Changed to Brightness temperature (BT) the observation in this 1231.3 cm^{-1} channel (BT1231) therefore serves as a measure for the cloudiness of an observation : if there are no or low or optically thin clouds, it will effectively measure the surface temperature, but as the clouds get thicker and higher, it will measure the cold cloud top temperatures. For any tile during any 16 day observation periods, we ~~can~~ compute quantiles Q based on the observed BT1231 to ~~screen design a filter that chooses~~ between cloudy and partially clear scenes ~~—We chose for every tile. We describe below the testing of the~~ different BT1231 quantiles ~~(so where~~ quantile $Q_{0.XY}$ will have a numerical value $BT_{1231,Q_{0.XY}}$ associated with it) ~~and show below the data contained between $Q_{0.90}$ and $Q_{1.00}$ can be considered “almost free of clouds.” to determine which value best provides nominally clear scenes for every tile (over ocean and land) that agree with other nominally clear datasets we have used previously~~ (Strow & DeSouza-Machado, 2020).

Figure 1 shows all the BT1231 observations for a chosen 16 day timestep in the form of a zonally averaged histogram (normalized probability distribution functions (PDFs)), with latitude on the vertical axis and BT1231 on the horizontal axis. The colorbar is the PDF value, and we used ~~data observations~~ spanning August 27, 2012 - September 11, 2012 which is approximately half way through the 20 year AIRS mission dataset used in this paper. The curves show the zonally averaged BT1231 values of the minimum ($Q_{0.00}$) in ~~dark-cyan~~blue, mean (thick red), median ($Q_{0.50}$ in orange), maximum ($Q_{1.00}$ in ~~light cyan~~) ; also shown are a handful of other zonally averaged BT1231 values, for example green) and $Q_{0.80}$, $Q_{0.90}$ (thick black curve); ~~We did not show other warmer quantiles such as $Q_{0.80}$ $Q_{0.95}$ and $Q_{0.97}$ —The since they are only slightly offset, either to the left (cooler) or right (warmer) as appropriate, relative to the $Q_{0.90}$ curve. The exception is that at the equator, $Q_{0.80}$ still has the remnants of lower temperatures due to clouds and is slightly cooler, as similarly seen in the behavior of the mean and median curves.~~




Figure 1. Zonally averaged BT1231 normalized histograms (probability distribution functions (pdf)) as a function of latitude and temperature bin, for ~~an~~ the 16 day timespan between 2012/08/27 - 2012/09/11. The vertical axis is in degrees Latitude and quantiles (the horizontal axis units are in Kelvin, while the colorbar units for the pdfs are in normalized counts per Kelvin). We also plot quantile curves $Q_{0.XY}$ which stand for the actual numerical value of the BT1231 quantile, as explained in the text. The thick black curve is the $Q_{0.90}$ quantile (and above) used in this paper, and is very close to the maximum $Q_{1.00}$ quantile. For clarity we have not shown other “warmer” quantiles such as $Q_{0.80}$, $Q_{0.95}$ since they are offset very close to the left and right of $Q_{0.90}$ respectively. The 210 K cutoff means we do not show the tail of the distribution of the observations over the winter polar regions, or the extremely cold DCC in the tropics.

The distributions are skewed to the left (negative skewness), as confirmed by the mean being less than the median. We also point out that even $Q_{0.80}$ sees much of the surface from the southern tropics to the northern polar region. The 220 K horizontal axis cutoff means we do not see the very cold (190 K) observations over the winter Antarctic.

The figure shows the expected qualitative features, for example (1) the tropical PDFs peak at around 295 K, but show some warmer observations, as well much colder observations (below 230 K) corresponding to Deep Convective Clouds (DCC); this gives a dynamic range of almost 100 K at the tropics (2) the BT1231 observed over the Southern Polar (polar winter) regions are much colder than the BT1231 observed over the Northern Polar (polar summer) regions and (3) the reddish peaks in the 30°N - 40°N are a combination of the marine boundary layer (MBL) clouds and warmer summer land temperatures. Figure 1 shows on average the cloud effect at the tropics is an additional modest 20 K (difference between $Q_{0.90}$ and $Q_{0.50}$) compared to the 100 K dynamic range. This is because the cloud fractions and cloud decks in the individual observations have effectively more clouds (with larger cloud fraction in the FOV) lower in the atmosphere than higher up; the net effect is that in the window region the atmosphere is on average radiating from the lower (warmer) altitudes, and so spectra whose BT1231 values are larger than $BT1231_{Q_{0.80}}$, see much of the surface emission as well.

547

548 We now use the above plots to select “almost clear” scenes. For any one tile, we de-
 549 fine set $\Psi_{0.XY}$ to have all observations i whose BT1231 lies between quantiles $Q_{0.XY}$
 550 and $Q_{1.00}$, $\{i \mid BT1231_{Q_{0.XY}} \leq BT1231(i) \leq BT1231_{Q_{1.00}}\}$. In what follows $Q_{0.XY}$
 551 is the radiances averaged over all the observations i which are in the set $\Psi_{0.XY}$, namely

$$r_{Q_{0.XY}}(\nu) = \frac{1}{N_{0.XY}} \sum_{i \in \Psi_{0.XY}} r_i(\nu) \quad (1)$$

552 where $r_i(\nu)$ are the $N_{0.XY}$ individual observations in set $\Psi_{0.XY}$. In this section we only
 553 use the $\nu = 1231 \text{ cm}^{-1}$ channel, but in later sections we easily form averages for all 2645
 554 channels, at any 16 day time step for any tile.

555 We tested different quantile sets $\Psi_{0.XY}$ to see which one can reliably be considered
 556 to provide a nominally “cloud free” global [observational](#) dataset, and chose the $Q_{0.90}$ av-
 557 erage (ie defined as averaged over the $\Psi_{0.90}$ set, [which spans \$Q_{0.90}\$ to \$Q_{1.00}\$](#)) as the one
 558 to use for the rest of this paper, unless explicitly stated otherwise. The tests primarily
 559 involved comparisons to scenes produced by the uniform/clear sky filter described in (Strow
 560 & DeSouza-Machado, 2020) for the same August 27, 2012 - September 11, 2012 sixteen
 561 day timespan. This latter filter selects clear scenes by both testing for uniformity (to within
 562 0.5 K) across a 3×3 grouping of AIRS scenes and also using a criteria that the observed
 563 window channel observations should be within ± 4 K of clear-sky simulations using ther-
 564 modynamic parameters supplied by reanalysis models. The results are shown in the left
 565 hand plot of Figure 2, plotted on a $1^\circ \times 1^\circ$ grid. We note in this plot the uniform/clear
 566 scenes that are plotted are limited to those over ocean, and for solar zenith less than 90°
 567 (daytime), which automatically filtered out many of the views over the (wintertime) South-
 568 ern Polar region. Immediately apparent are the gaps produced by the uniform/clear fil-
 569 ter *e.g.* in the Tropical West Pacific or off the western coasts of continents where there
 570 are clouds. The gaps can be changed by *e.g.* changing the 4K threshold to allow more
 571 or fewer scenes through the filter.

572 The center plot shows for all tiles, the daytime scenes selected by the $Q_{0.90}$ filter
 573 for the same time period, on the same $1^\circ \times 1^\circ$ grid. Compared to the left hand plot, the
 574 spatial coverage is almost complete, as the $Q_{0.90}$ average always has the hottest 10% of
 575 the observations. At this 1° resolution, used for comparison with the uniform/clear grid
 576 filter described in the previous paragraph, gaps are seen in regions where for example
 577 the local topography means observations over mountains would be colder than the sur-
 578 rounding coastal or plain regions. This is not a concern since zooming back out to the
 579 coarser $3^\circ \times 5^\circ$ tile resolution, will include $Q_{0.90}$ [data-observations](#) for the quantile and
 580 trending analysis.

581 To compare the mean observations we remove the over-land and over-polar region
 582 [data-observations](#) from the center plot. The right hand plot shows the mean observed
 583 BT1231 from the $1^\circ \times 1^\circ$ grid from the uniform/clear sky filter as a function of latitude,
 584 compared to the $1^\circ \times 1^\circ$ grid from the $Q_{0.90}$ scenes. The difference between the unifor-
 585 m/clear versus $Q_{0.90}$ average is within about $0.25 \text{ K} \pm 1 \text{ K}$ across the southern trop-
 586 ics to the northern midlatitudes, though the bias rises to about 1 K by about -50°S . We
 587 consider this an acceptable difference, as we could tune the thresholds for the uniform/-
 588 clear filter to *e.g.* change the areal coverage and/or number of clear scenes and hence
 589 comparisons to the $Q_{0.90}$ scenes.

590

591 The results presented in this section have been checked for robustness, using other
 592 16 day intervals spanning the four seasons. We conclude that for any 16 day timestep




Figure 2. Clear scenes for the same 2012/08/27 - 2012/09/11 timespan selected by (left) an uniform/clear sky filter and (center) the Q0.90 BT1231 average described in this paper. The colorbars for the left and center plots are in Kelvin. The right hand plot shows the mean (over ocean) observed BT1231 (vertical axis, in Kelvin) as a function of latitude, for the two selections; the difference is about $0 \text{ K} \pm 1 \text{ K}$ in most region-regions except in the southern midlatitudes where the Q0.90 average produced scenes that were about 1 K cooler on average. Note that in this and subsequent figures, Q0.90 is the average of all data points values between Q0.90 (shown in Figure 1) and the maximum, using observed BT1231 as the discriminator as explained in the text.

the radiances used in the Q0.90 average (a) produces almost complete spatial coverage of the Earth, (b) selects scenes whose average BT1231 is very close to the average BT1231 from scenes selected using an uniform/clear filter (c) trends from that quantile typically differ by less than $\pm 0.002 \text{ K yr}^{-1}$ from the other quantiles and (d) this selection produces spectral trends which compare well against those obtained from the quality assured binned AIRS CCR data record (Manning, 2022), and reinforces the notion that our quantile based selection is selecting nominally clear scenes. Together these imply the Q0.90 average is an acceptable proxy for “clear scenes”. For the remainder of the paper we therefore consider Q0.90 as consisting of nominally clear observations whose BT1231 lies between the 90th quantile and hottest observation. Our retrievals using this $\text{Q0.90} \rightarrow \text{Q1.00}$ averaged observational dataset (shortened to Q0.90) is referred to as AIRS_RT in what follows.

3.2 Observed trends from the Q0.90 Quantiles

Having selected the Q0.90 observations, for each tile the average radiance per 16 day interval is computed. With two sixteen day periods not available (Aqua platform or AIRS shutdowns during *e.g.* solar flare events) this gives a total of 457 time steps over 20 years. Anomalies are formed from this time series, and then de-seasonalized to give the spectral radiance trends and error estimates (Strow & DeSouza-Machado, 2020) using Matlab *robustfit*:

$$r_{\text{observations}}^{16 \text{ days}}(t) \sim r_{\text{fit}}(t) = r_o + a_1 t + \sum_{i=1}^4 c_i \sin(n2\pi t + \phi_i) \quad (2)$$

with a_1 and its associated uncertainty, both converted to brightness temperature (BT), being the trends in K yr^{-1} . Using sub-harmonics in the fit did not produce any noticeable change in the AIRS_RT retrievals (described below).

The left panel of Figure 3 shows the descending orbit (nighttime) 20 year (September 2002- August 2022) global averaged spectral observations for the five quantiles mentioned above. We note the spectra in most of the plots in this section are weighted by the $\cosine(\text{latitude})$ of the tiles, unless otherwise stated. In addition we only show the

619 640-1640 cm^{-1} region, and ignore the shortwave 2050-2750 cm^{-1} region since the AIRS
 620 SW channels are drifting relative to the LW (Strow & DeSouza-Machado, 2020). Spec-
 621 tral averages constructed from Figure 1 would have this same behavior, namely that in
 622 the window region the mean spectrum of ~~data-observations~~ populating the warmer quan-
 623 tiles (Q0.80, Q0.90, Q0.95, Q0.97) as defined in Equation 1 are on the order of a Kelvin
 624 apart, and have about half/quarter that difference in the optically thicker regions dom-
 625 inated by H_2O and/or CO_2 absorption respectively.

626 The right hand panel of Figure 3 shows (top) the trends and (bottom) the 2σ trend
 627 uncertainties for these quantiles, in K yr^{-1} . We emphasize that the top right panel shows
 628 that the spectral trends for the quantiles lie almost on top of each other; the difference
 629 between the Q0.50 and other trends is at most about $+0.003 \text{ K yr}^{-1}$ (out of a 0.02 K
 630 yr^{-1} signal) in the window region (and about $+0.0045 \text{ K yr}^{-1}$ in the troposphere tem-
 631 perature sounding channels), or less than 10%. Similarly the largest trend uncertainty
 632 in the bottom panel is for Q0.50. This implies that clouds effects in the infrared ~~do~~ pro-
 633 duce the largest variability (blue curve) ~~but~~. Globally on average for the infrared ~~are not~~
 634 ~~changing much, so the spectral trends for all quantiles, ranging from clearest (Q0.97) to~~
 635 ~~allsky (Q0.50 very similar, but differences are seen on regional scales. This implies the~~
 636 $+0.022 \text{ K yr}^{-1}$ window region trends are dominated by surface temperatures changes
 637 and to a lesser extent by water vapor changes.”

638 ~~TOA radiances in the 15- μm (700-800 cm^{-1}) region are impacted by two effects (X. Huang et al. (2023); Raghura-~~
 639 ~~and our work all show, either in radiance or OLR space, (a) the increased optical depths~~
 640 ~~due to increasing atmospheric-observed radiance in the window channels, due to surface~~
 641 ~~temperature increases (b) the $\approx -0.06 \text{ K yr}^{-1}$ decrease in BT in the 700-750 cm^{-1} troposphere~~
 642 ~~sounding region, which is due to a combination of the CO_2 leads amounts/optical depth~~
 643 ~~risers leading to atmospheric emission from higher altitudes/lower temperatures, resulting~~
 644 ~~in almost a -0.06 K/year signal for the troposphere, and (b) the together with atmospheric~~
 645 ~~temperature increases (again about shown later in this paper to be between $+0.01$ to $+0.02$~~
 646 ~~K yr^{-1}); (c) increases in the 1350-1640 cm^{-1} free troposphere water vapor sounding region~~
 647 ~~and (d) the 1280-1340 cm^{-1} decreases are due to CH_4 increases.~~

648 Also of interest ~~is are~~ the trends in the stratosphere (650-700 cm^{-1}) changes which
 649 consists of a stratospheric cooling signal (negative) and emission higher up due to increased
 650 CO_2 ; combining to give a net zero effect over 20 years, also seen in (Raghuraman et al.,
 651 2023). The H_2O signal is evident in the 1400-1625 cm^{-1} region, and is only slightly pos-
 652 itive; in other words, increasing temperatures have led to increased atmospheric amounts
 653 of H_2O , and the water vapor feedback has reduced the amount of outgoing flux in that
 654 region. By extension, this can ~~only also~~ be expected to have happened in Far Infrared
 655 (10-650 cm^{-1}) spectral regions affected by water vapor, ~~and but~~ cannot be wholly con-
 656 firmed as current sounders do not make direct measurements in that region. In the near
 657 future it is anticipated the Far Infrared Outgoing Radiation Understanding and Mon-
 658 itoring (FORUM) mission (Palchetti et al., 2020) will provide ~~data-observations~~ to fill
 659 in this important observation gap.

660 ~~20-year trends from different observation quantiles. The left hand panel shows the~~
 661 ~~mean globally averaged BT observations from 20 years of AIRS data, for quantiles Q0.50,0.80,0.90,0.95,0.97~~
 662 ~~as described in the text. The right hand panel shows (top) the globally averaged trends~~
 663 ~~for those different quantiles and (bottom) the spectral uncertainty in the trends. The~~
 664 ~~nighttime (descending) trends are shown in these plots.~~

665 4 ~~Spectral closure : comparisons between observed and simulated spectral~~ 666 ~~trends~~

667 Previous work has demonstrated that the radiances from AIRS are climate quality,
 668 if one restricts the channel set to the ~ 450 channel set that is largely immune to nonphysical


drifts. In closing this section we describe a way to test the quality of the monthly thermodynamic output from reanalysis and point out a comparison of spectral trends between AIRS_L3 observations and reanalysis/or L3 products which are all in geophysical space, against the AIRS L1C observational data which is in radiance space. This is accomplished by geolocating the monthly (ERA5) surface temperature, air temperature, water vapor and ozone fields to tile centers as described in Section 2.2, which are then input and run through the fast model, for the entire 20 years. Spectral radiance trends were then computed from these time series of (clear sky) spectral radiances. The conversion of L3 retrieval and NWP reanalysis trends to a radiance time series, provides a rigorous check of their accuracy against the observed AIRS L1C radiance trends which are validated to be highly accurate.

The simulations included realistic column linearly increasing with time mixing ratios for L3 simulations is presented and discussed in Figures B1 and for the ERA5 spectra, as well as land or ocean surface emissivity co-located to tile centers together with view angles of about 22, which is the average view angle of the tiled observations. From these the ERA5 spectral trends were derived similarly to what was described above for the AIRS observation spectral trends. B2 of Appendix B.

4 Testing the variability of representative points from NWP reanalysis

Each sixteen day $3^\circ \times 5^\circ$ tile contains ~ 12000 observations, which means for each tile about 600 daytime and 600 nighttime observations are averaged to produce the Q0.90 observational dataset per timestep. Conversely there are typically only ~ 240 monthly ERA5 0.25° points per $3^\circ \times 5^\circ$ tile; for 1° resolution AIRS L3 and CLIMCAPS L3 there are even fewer (15) points per tile. This low number of points means we chose a simple solution of using the grid cell closest to the center of each $3^\circ \times 5^\circ$ tile for building the NWP reanalysis and L3 geophysical time series. This choice is validated below using the following test to see for example how surface temperature trends would be impacted as we changed the representative point for the ERA5 model fields.

For the descending overpass we built complete sets of approximately 240 ERA5 points per tile per month; at 0.25° resolution one of these is almost certainly at the tile center. From these monthly sets, we could either directly read the tile center temperature (our default), or compute the average surface temperature per tile, or compute the average of the hottest 10% surface temperatures per tile. This was done for all 20 years (240 monthly timesteps) after which the three timeseries were trended. Over ocean the differences between all three datasets/sets of data was typically $-0.001 \pm 0.005 \text{ K yr}^{-1}$, while over land the differences were about $0.001 \pm 0.01 \text{ K yr}^{-1}$. This is to be compared to mean trends of about $0.014 \pm 0.02 \text{ K yr}^{-1}$ over ocean and $0.025 \pm 0.04 \text{ K yr}^{-1}$ over land: the spread of the ocean and land ERA5 surface temperature trends for the three methods, is much smaller than the mean trends. Given that there were far fewer re-analysis points in a grid box than tiled Q0.90 observations, coupled with the fact that choosing the 10% warmest profiles would provide an even smaller sample, we chose to use the tile center to be the representative point to co-locate the model fields.



20-year zonally-averaged spectral brightness temperature trends (in K yr^{-1}) for (left) AIRS Q0.90 observations and (right) clear sky simulations using ERA5 monthly model fields. The center panel shows the AIRS Q0.90 spectral uncertainties. The ERA5 simulations included linear trends of $\frac{1}{1000}$ and $\frac{1}{1000}$, while the trends in ERA5 are from the reanalysis itself.

Figure B2 shows the AIRS observed Q0.90 (nominally clear) descending (night) zonally averaged results in K yr^{-1} in the left panel, and the zonally averaged simulated clearsky (without clouds) spectral trends (also in K yr^{-1}) from monthly ERA5 fields in the right panel. The center panel shows the spectral trend uncertainties from the observations, also in K yr^{-1} . In the next section we derive geophysical trends from these (AIRS observed) spectral trends, and the similarities/ differences in geophysical trends between observations and models/operational data can be partially understood from the similarities/differences in the spectral trends. For example, the sounding region (1350-1600) of the left and right panels of Figure B2 shows roughly similar (positive) spectral trends in the tropics and mid-latitudes; there are some slight differences in the high altitude channels (1450-1550 region). The following sections shows that this results in subtle differences in the tropospheric water vapor trends. Observations and simulations both have positive $\text{dB T}/\text{dt}$ in the 800-960, 1150-1250 region, indicating surface warming; however the ERA5 simulation show more warming in the southern polar regions than do the AIRS observations. Note the mean warming in the tropics for both observations and ERA5 simulations is less than that in the mid-latitudes, and the polar regions show the largest overall change in brightness temperature in the window region. Large differences are seen in the 10 μm (1000) sounding region, which are not surprising since ozone assimilation is not a primary goal of ECMWF assimilation; here we do not address these as we focus on the changes to the moist thermodynamic state. The window region trends computed using the ERA5 model are more positive in the Southern Polar region. Conversely the 640-700 spectral region is positive, especially in the tropics; however the observations show a net cooling trend away from the tropics, compared to the ERA simulations. This demonstrates the importance of the model \rightarrow spectral trend comparisons, given the accuracy of the AIRS observations.

The paper by Raghuraman et al. (2023) shows similar figures, but in terms of spectral OLR trends encompassing the 0-2000 range, while X. Huang et al. (2023) shows similar plots for a slightly smaller time period (2002-2020) using the nadir L1B radiance dataset which has no or minimal frequency corrections compared to the L1C set we use in this paper. X. Huang et al. (2023); Raghuraman et al. (2023) and our work all show, either in radiance or OLR space, (a) the increased observed radiance in the window channels, due to surface temperature increases (b) the $\simeq -0.06 \text{ K yr}^{-1}$ decrease in BT in the 700-750 troposphere sounding region, which is due to the amounts increasing; we also see differences in the signs of the BT changes in the 650-700 stratospheric and temperature channels for some latitudes between observations and ERA5 simulations (c) increases in the 1350-1640 water vapor sounding region seen in Figures 3 and B1, and (d) the 1280-1340 decreases are due to increases.

3.1 Sample spectral closure comparisons using other monthly products

Here we follow the earlier work of and convert the ERA5 monthly model fields to spectral radiances, after which we compute spectral trends for comparison to AIRS observations. Spectral closure calculations for the entire 20-year timeseries were also generated for the monthly MERRA2 model fields, as well as the monthly AIRS v7 L3 and CLIMCAPS L3 retrieved data products. Again only the monthly thermodynamics and surface temperature fields for all 72×64 tiles were used in these runs, with GHG changes added in for each timestep as described above. Spectral trends were then computed using Equation 2.

We chose just one limited example here to illustrate the power of this approach for diagnosing which dataset is more accurate, given that the AIRS spectral trend accuracy

5 Geophysical Trend Retrieval outline

5.1 Setting up the Retrieval Problem

The observed clear sky spectral brightness temperature for a tile at any time t can be modeled as

$$BT(\nu, t) = f(X(t), \epsilon(\nu, t), \theta(t)) + \text{NeDT}_{\text{retrieval}}(\nu) \quad (3)$$

where the state vector $X(t)$ has the following five geophysical state parameters : (1) surface temperature (ST), (2) atmospheric temperature profile $T(z)$, (3) water vapor profile $WV(z)$, (4) ozone profile $O3(z)$ (5) greenhouse gas forcings (GHG) due to CO_2 , CH_4 and N_2O changing as a function of time t and $f(X(t), \epsilon, \theta, \nu)$ is the clear sky radiative transfer equation for channel center frequency ν . The spectral noise ~~NeDT(ν) for~~ NeDT_{retrieval}(ν) varies with scene temperatures and on particulars of the retrieval algorithm. For single footprint retrievals using daily observations, the spectral noise NeDT_{retrieval}(ν) in a typical tropical “clear scene” is about 0.1 K in window region, increasing to about 1 K in the 15 μm temperature sounding channels and about 0.2 K in the 6.7 μm water vapor sounding region, but the noise will vary as a function of scene temperature and is usually larger for operational L2 retrievals which use cloud clearing. We parametrize the GHGs using single numbers (such as ppm(t) for the CO_2 column), and include the AIRS orbit and viewing angle geometry θ and the surface emissivity $\epsilon(\nu)$, while we omit forward model and spectroscopy errors. We ignore cloud scattering as well as the spatial variation of the state parameters, emissivity and scan angle geometry within a tile. Linearizing the above equation about the time averaged profile, the relationship between the observed spectral trends and desired thermodynamic trends is given by

$$\frac{dBT(\nu)}{dt} = \frac{\partial f}{\partial \bar{X}} \frac{d}{dt} \bar{X}(t) = K(\nu) \frac{d}{dt} \bar{X}(t) + K_{\text{emissivity}}(\nu) \frac{d}{dt} \epsilon(t) \xrightarrow{0} K(\nu) \frac{d}{dt} \bar{X}(t) \quad (4)$$

where the matrix $K(\nu)$ is the thermodynamic jacobian (surface temperature, air temperature and trace gases) and we ignore any orbit drifts (changes to θ), instrument changes (changes to ~~NeDT(ν)~~ NeDT_{retrieval}(ν)) and surface emissivity ($\epsilon(\nu)$); the last assumption is investigated in a later section. The overbars on parameters X denotes this is a time average (linear trend) that we are working with, and we have converted from radiances in Equation 2 to brightness temperatures in Equations 3 and 4.

5.2 Jacobian calculations

For a typical clear sky tropical sky atmosphere, the 800 - 1200 cm^{-1} window region has surface temperature (SKT) jacobians which are about +0.5 to +0.75 K per degree SKT change and -0.75 to -0.25 K per 10% change in column water vapor. The spectral variability in these window region jacobians is primarily due to reducing water continuum absorption as you move from the 800 cm^{-1} end to the 1200 cm^{-1} ; consequently the surface temperature jacobians becomes closer to unity and the column water jacobians become closer to zero as water vapor amount decreases (drier atmospheres in the mid-latitudes and polar regions). The hyperspectral channels used in this work assist in partitioning these two competing changes (though not perfectly), which we validate against other datasets in this study. As seen in Figure B2 typical magnitudes of the spectral trends on the left hand side of Equation 4 are less than about 0.1 K per year. Equation 4 is in the usual inversion form $\delta y = K \delta x$, and the Optimal Estimation Rodgers (2000) solution used to solve the anomaly time series in (Strow et al., 2021) is also used here. The noise term ~~used-NeDT_{retrieval}(ν) for the trend retrieval-NeDT(ν) is not the instrument noise since each 16 day point in our time series is averaged over hundreds of observations as earlier described; instead the uncertainty is that due to retrievals is now the uncertainty that naturally arises from the~~ inter-annual variability ~~in the linear trends obtained from the trend fitting when doing the linear trend fitting shown~~ in Equation 2. Examples of typical noise values are shown in the bottom right hand panel of Figure 3.

ERA5 monthly model fields at tile centers, together with time varying concentrations of GHG such as CO₂, were averaged over 20 years so jacobians could be computed. The GHG concentrations were a latitude dependent increase of about 2.2 ppm yr⁻¹ for CO₂ derived from the CarbonTracker (Peters et al., 2007) (CarbonTracker CT-NRT.v2023-4, <http://carbontracker.noaa.gov>) ~~model~~-data at 500 mb. Our pseudo-monochromatic line by line code kCARTA (De Souza-Machado et al., 2018, 2020) was used with these averaged profiles to produce accurate analytic jacobians. The HITRAN 2020 line parameter database (Gordon & Rothman, 2022), together with MT-CKD 3.2 and CO₂,CH₄ line mixing from the LBLRTM suite of models (Clough et al., 2005) were used in the kCARTA optical depth database (De Souza-Machado et al., 2018). A 12 month geographical land-varying spectral emissivity database spanning one year from (Zhou et al., 2011) was used, while ocean emissivity came from (Masuda et al., 1988). The atmospheric temperature, water vapor and ozone profile jacobians, and the surface temperature and column jacobians for the GHG gases such as CO₂ and CH₄ and N₂O, were then convolved using the best estimate AIRS Spectral Response Functions (Strow, Hannon, Weiler, et al., 2003).

Tests done for this paper, together with the results in (Strow et al., 2021), established that jacobians derived from MERRA2 versus ERA5 produced no significant differences in the context of retrieved trends or anomalies done for this paper, as the uncertainty in linear trends due to inter-annual variability dominates over any uncertainty (or differences between) model fields.

5.3 Optimal Estimation Retrieval : State vector, covariance matrices and *a-priori*

Using monthly ERA5 model fields averaged over 20 years, for each of the 64 × 72 tiles we computed analytic jacobians for the following (vector) atmospheric thermodynamic variables [fractional water vapor, fractional ozone and temperature] together with (scalar) surface temperature. We retrieved fractional gas concentration trends $dfrac{X}/dt = 1/X_{avg}(z)dX_{avg}(z)/dt$ to keep all values in the state vector at about the same magnitude. A single iteration Optimal Estimation retrieval (Rodgers, 2000) is used to simultaneously solve for the geophysical parameter trends. As in Strow & DeSouza-Machado (2020) the geophysical covariance uncertainty matrices are a combination of Tikonov and covariance regularization. The uncertainties for the covariance matrices were typically [0.1,0.25,0.45] K yr⁻¹ for the surface/tropospheric/stratospheric temperature trends, and [0.04/0.02] yr⁻¹ for the fractional tropospheric/stratospheric water vapor trends. Tikonov L1 regularization Rodgers (2000) also included, with the scalar factor multiplying this regularization corresponding to about 1/10 the covariance uncertainties. The spectral uncertainties used in the retrievals come from the above mentioned trend uncertainties. For completeness we note that a sequential ~~retrieval (see for example)~~ trend retrieval produces very similar geophysical trends.

Here we emphasize four unique points about our geophysical trend retrievals, which ~~sets us apart~~ distinguishes this approach from trends derived from other datasets. Firstly the *a-priori* trend state vector is zero ($dST/dt = dT(z)/dt = dQ(z)/dt = 0$) for all geophysical parameters, except for water vapor where we enforced constant (or slightly increasing) relative humidity as described below. This ensures traceability of our retrieval is straightforward especially wherever the AIRS instrument has sensitivity. For example the 300 - 800 mb water vapor trend retrievals will be based on the observed data only, thereby insulating us from any possible *a-priori* information from *e.g.* climatology or ~~NWP models~~ reanalysis, unlike the operational AIRS V7 or CLIMCAPS retrievals which use first guesses based on neural net and MERRA2 respectively.

Secondly ~~as seen in Figures B2 and B1, in~~ the 15 μm region ~~there is of Figure B2~~ shows a large spectral overlap signal (-0.06 K yr⁻¹) from the increasing CO₂, which is much larger than the expected atmospheric temperature trend (0.01 - 0.02 K yr⁻¹). These

correlations makes it difficult to jointly retrieve both temperatures changes and changes in well mixed GHGs such as CO₂. We chose to focus on retrieving temperature changes only, by spectrally removing the effects of changing CO₂, CH₄ and N₂O GHG concentrations. This was done by using the GHG trends estimated from NOAA ESRL Carbon-Tracker data multiplied by the appropriate GHG gas column jacobian (CO₂, N₂O and CH₄ and CFC11, CFC12) computed as described above using the averaged over 20 years ERA5 monthly profile for each tile.

Thirdly instead of using all 100 layers described in the AIRS forward model (Strow, Hannon, DeSouza-Machado, et al., 2003), we combine pairs of layers for a 50 atmospheric layer retrieval, as the AIRS radiances contain far fewer than 100 pieces of information (see *e.g.* (Maddy & Barnett, 2008; De Souza-Machado et al., 2018)).

Fourthly, modern hyperspectral infrared sounders have highest sensitivity to temperature and water vapor in the mid-tropopause; see for example the averaging kernels in (Irion et al., 2018). Using a zero fractional WV trends *a-priori* at all levels, it was fairly straightforward to obtain fractional WV(z) trends close to those from the ~~NWP-model~~ reanalysis datasets in the 300-850 mb region. In order to improve our results in the lowest layers, we enforced a constant relative humidity approximation, which is a well-known, expected behavior under global climate change (Soden & Held, 2006; Sherwood et al., 2010). This was done by ignoring the contribution due to water vapor changes in the observed BT1231 trend, and using it as an approximation for air temperature trend over ocean; this allows us to compute an estimate of how the water vapor would need to change

$$RH(T) = \frac{e}{e_{sat}(T)} \implies \delta(RH) = \frac{1}{e_{sat}(T)} \delta e - \frac{e}{e_{sat}^2(T)} \delta e_{sat}(T) = \frac{1}{e_{sat}(T)} \delta e - \frac{e}{e_{sat}(T)} \frac{L_v}{R_v} \frac{1}{T^2} \delta T \quad (5)$$

where $e, e_{sat}(T)$ are the vapor pressures and we used $e_{sat}(T) = e_{s0} e^{\frac{L_v}{R_v} (\frac{1}{T_0} - \frac{1}{T})}$ (where L_v, R_v are latent heat of vaporization and gas constant respectively) to go from the expression in the center to the expression on the right. If we expect the change in RH to be zero then $\frac{\delta e}{e} = \frac{L_v}{R_v} \frac{\delta T}{T^2}$, where we can use $\delta T / \delta t \sim d/dt BT1231$. to approximate the *a-priori* fractional vapor pressure rates (or *a-priori* fractional water vapor rates) between surface and 850 mb, smoothly tailing to 0 in the upper atmosphere. Subsection 6.2 has a similar discussion on a proposed method to alleviate the lack of sensitivity to upper atmosphere water vapor. Our default results in this paper are from using the MLS *a-priori*, unless otherwise stated.

5.4 Testing on synthetic trend spectra made from ERA5 Reanalysis monthly fields

We tested the retrieval code by using it on the simulated nighttime only ERA5 spectral trends, and compared to geophysical trends computed directly from the ERA5 reanalysis~~model~~. Spot checks of the spatial correlations of ERA5 fractional water vapor and temperature trends versus the trends retrieved from synthetic spectra/our retrieval algorithm, peaked at 500 mb with correlations of about 0.9, compared to 800 mb correlations of 0.80 and 0.55 for temperature and fractional water vapor trends respectively and 200 mb correlations of 0.89 and 0.69 for dT/dt, dWVfrac/dt. This is to be expected since a computation of the water vapor averaging kernels for infrared instruments for arbitrary atmospheric profiles typically shows they peak in the 300 mb - 850 mb range and decrease rapidly away from those regions; conversely the temperature averaging kernels stay relatively uniform through the free troposphere and above, though they also decrease close to the surface; see for example (Irion et al., 2018; Smith & Barnett, 2020; Wu et al., 2023).

Figure 4 shows a sample set of results using nighttime ERA5 model output converted to spectral trends as described above. The top panels (A) are always the atmospheric trends computed directly from the monthly ERA5 model fields, while the bottom pan-




Figure 4. Comparing geophysical trends derived directly from ERA5 monthly nighttime fields (top) vs from the AIRS_RT retrieval applied to the ERA5 reconstructed spectral trends (bottom). Left panel is dT/dt (Horizontal axis are all in $K\text{-yr}^{-1}$ latitude (deg)) while rightmost panel vertical axis is $d\text{fracWV}/dt$ (Note the vertical axis is logarithmic for the temperature trends and linear for the water vapor trends. The colorbar in for the left panels is $K\text{-yr}^{-1}$ while the colorbar for the right panels is yr^{-1} (as fractional water vapor has no units)).

els (B) are the atmospheric trends retrieved from the converted ERA5 spectral brightness temperature trends. The left most panel is the atmospheric temperature trend comparison (both in $K\text{-yr}^{-1}$) while the rightmost panel is the fractional atmospheric water vapor trend comparison (in yr^{-1}).

It is evident from the figure that the tropospheric trends in the tropical and mid-latitude regions are quite similar, and there are differences in the polar regions and stratospheric regions where the AIRS instrument has reduced sensitivity. The atmospheric and surface trends are shown in Table 1, divided into “all” (which is the entire ± 90 latitude range and 0-1000 mb vertical range) and “T/M” which is the tropical/midlatitude region, which is further reduced to 050-900 mb for air temperature and 300-800 mb for water vapor. “ERA5 direct” are trends computed directly from the geophysical fields, while “ERA5 spectral” are retrieved from the spectral trends.

	dT_z/dt $K\text{-yr}^{-1}$ A SFC-TOA	dT_z/dt $K\text{-yr}^{-1}$ T/M 050-900 mb	$dSKT/dt$ $K\text{-yr}^{-1}$ A	$dSKT/dt$ $K\text{-yr}^{-1}$ T/M	$d\text{fracWV}/dt$ $K\text{-yr}^{-1}$ A GND-TOA	$d\text{fracWV}/dt$ $K\text{-yr}^{-1}$ T/M 300-800 mb
ERA5 direct	0.010 ± 0.038	0.029 ± 0.013	0.020 ± 0.035	0.018 ± 0.032	0.003 ± 0.002	0.002 ± 0.001
ERA5 spectral	0.004 ± 0.033	0.027 ± 0.012	0.019 ± 0.033	0.016 ± 0.029	0.001 ± 0.001	0.002 ± 0.001

Table 1. Cosine weighted air temperature and skin temperature trends (in $K\text{-yr}^{-1}$), and fractional water vapor trends (in yr^{-1}), together with uncertainties. The “ERA5 direct” are directly from the ERA5 geophysical trends, while “ERA5 spectral” are trends retrieved from the converted ERA5 spectral trends.

5.5 Surface emissivity changes

Equation 3 explicitly includes the surface emissivity in the equation of radiative transfer; however Equation 4 assumes this is unchanging. Here we rewrite Equation 4 as

$$\frac{d\overline{BT}(\nu)}{dt} - K_{emissivity}(\nu) \frac{d}{dt} \overline{\epsilon(t)} \rightarrow \frac{d\overline{BT}(\nu)}{dt} = K(\nu) \frac{d}{dt} \overline{X(t)} \quad (6)$$

Ocean emissivity has a dependence on windspeed (Masuda et al., 1988). (Lin & Oey, 2020) and other literature suggest wind speed increases of $+2.5 \text{ cm s}^{-1} \text{ yr}^{-1}$ have occurred between 1993-2015 in the tropical Pacific, and smaller (or close to zero) values elsewhere. The monthly ERA5 $u10, v10$ 10 m speeds for the 20 year time period in this paper also showed the maximum absolute trend was 0.09 m/s/year (over the Southern Ocean) while the global ocean mean and standard deviation were $0.006 \pm 0.022 \text{ m s}^{-1} \text{ yr}^{-1}$; The emissivity changes over ocean using a 0.025 m s^{-1} wind speed change are on average on the order of 1×10^{-6} per year in the thermal infrared window (or about 0.0003 K yr^{-1} change in the window region); assuming the optical properties of water do not substantially change with the $\sim 0.02 \text{ K}$ increases seen in all the datasets considered in this paper, these very small emissivity changes [due to windspeed changes](#) are of no consequence.

[We also estimate how the changing ocean temperatures would change the emissivity. Assuming no atmosphere, the radiance measured at the TOA is \$r_0\(\nu\) = \epsilon\(\nu\)B\(\nu, T_0\)\$ where \$T_0\$ is the temperature, \$\epsilon\$ is the emissivity and \$B\(\nu, T_0\)\$ is the Planck function. If the temperature is perturbed by \$\delta T\$ then the radiances changes by an amount \$\delta r\(\nu, T_0\) = \epsilon\(\nu\) \frac{dB\(\nu, T_0\)}{dT} \delta T + B\(\nu, T_0\) \frac{d\epsilon\(\nu, T_0\)}{dT} \delta T\$. The derivative of the Planck function is easily computed analytically. An estimate of the ocean emissivity change with temperature is \$\sim 2 \times 10^{-4}\$ per Kelvin, using the information in \(Newman et al., 2005; Nalli et al., 2022\). Inserting these numbers yields a BT change of \$\sim 1.5 \times 10^{-3} \text{ K}\$ due to the change in emissivity, which is much smaller than the assumed \$0.2 \text{ K}\$ ocean temperature change.](#)

Land emissivity changes were estimated as follows. A global monthly mean emissivity database, the Combined ASTER and MODIS Emissivity over Land (CAMEL v003) has recently been released (Borbas et al., 2018). We matched the tile centers to the database for the 20×12 months spanning our 2002/09 - 2022/08 time period, and computed the emissivity trends over land; the results (not shown here) were on the order of -1×10^{-4} and $+3 \times 10^{-4}$ in the $800\text{-}960 \text{ cm}^{-1}$ and $1100\text{-}1250 \text{ cm}^{-1}$ regions respectively, averaged over the land observations. For each tile the $K_{emissivity}(\nu) \frac{d}{dt} \overline{\epsilon(t)}$ term was estimated by running SARTA with the default emissivity, then differencing with the SARTA output obtained when the emissivity trends were added on. Averaged over the planet, the spectral changes arising from these emissivity changes were much smaller than the spectral trends seen in Figure 3, about -0.001 K yr^{-1} between $800\text{-}960 \text{ cm}^{-1}$ and about $+0.002 \text{ K yr}^{-1}$ on the $1100\text{-}1250 \text{ cm}^{-1}$ region (which we do not use in our retrieval, since many of the channels are synthetic and the real channels are drifting Strow et al. (2021)). The land only results were roughly three times these magnitudes. Using these emissivity jacobians on the left hand side of Equation 6 and running the retrieval on the adjusted spectral trends over land, resulted in about at most 0.01 K increases to the zonally averaged surface temperature changes over land; zonally averaged these largest differences were at about 40°N to 60°N and -25°S to $+15^\circ\text{N}$, due to emissivity decreases; the 20°N to $+35^\circ\text{N}$ region which included the Sahara and swathes of Asia, had emissivity increases but the averaged-over-land temperature decreases were small, as there were offsetting emissivity increases in other land areas at the same latitudes. We did not pursue the impact of these emissivity changes further as the CAMEL database is affected by the stability of the MODIS data, and our results below will not include accounting for changes in land emissivity.

6 Results

The trends retrieved in the previous section using simulated radiance trends show that the retrieval package is working as expected. Here we apply our retrieval to observed AIRS L1C radiance trends and compare the retrieved AIRS_RT geophysical trends to those computed directly from the ERA5/MERRA2 model fields and AIRS L3/CLIMCAPS L3 products. We will have an expectation that since the simulated radiance trends had no noise added to them, the uncertainty in the spectral rates was lower than the actual observed spectral uncertainty; this will lead to larger uncertainties and/or errors in our retrieval using observed radiance trends.

~~We will make most comparisons against NWP models.~~ Most of the comparisons against reanalysis model fields and L3 products will be made in the context of averages over the descending/night (N) and ascending/day (D) data-observations since the MERRA2 (and GISS) datasets are only available as a D/N-average; the reader is referred to the Appendix where we show a few of the D-N differences. The results are shown in the order of surface/column trends (surface temperature and column water), followed by zonal averages of the atmospheric temperature and fractional water vapor trends. We also refer the reader to Appendix B which presents an interpretation of these geophysical trend comparisons, using trends in radiance spectral space.

6.1 Skin Temperature trends

There are typically multiple (window) channels that are sensitive to a surface pressure, meaning the radiances typically have more information content for the surface temperature (assuming the surface emissivity is well known and there are no clouds) rather than for example air temperature. Figure 5 shows the diurnally averaged day/night (D/N) surface temperature trends from 6 datasets : AIRS_RT observations, AIRS L3, CLIMCAPS L3, ERA5, MERRA2 and NASA GISTEMP. AIRS_RT shows an overall global warming of $+0.021 \text{ K yr}^{-1}$; the cooling trends include the tropical eastern Pacific and south of Greenland and tropical northern Atlantic. The rest of the datasets also show similar patterns of cooling in the N. Atlantic Ocean, warming over the Arctic and some degree of cooling over the Antarctic Ice Shelf/Southern Ocean as does AIRS_RT. The AIRS v7 L3 shows some cooling over Central Africa and the Amazon not seen in the AIRS_RT trends, where one could expect Deep Convective Clouds and possible cloud clearing issues. We also point out the AIRS L3 product has many missing values off the western coasts of N. and S. America, due to cloud clearing issues. MERRA2 shows ~~more cooling~~ significant cooling trends over C. Africa, ~~and just like the AIRS v7 data, a lot of cooling~~ and near the Antarctic Ice Shelf. Of note here is that although CLIMCAPS uses MERRA2 as its first guess, their surface temperature trends are not similar, especially around the Antarctic where MERRA2 shows strong cooling trends. Over the ocean GISS shows similar trends to what AIRS_RT trends show. An earlier study of Land Surface Temperatures between 2003-2017 using MODIS (Prakash & Norouzi, 2020) shows very similar large daytime cooling trends over parts of central and western Indian subcontinent that we see from our retrieval as well as directly from the BT1231 channel trends; for tiles that straddle both ocean and land the quantile method picks up the hottest observations, which especially during summer are mostly over the Indian subcontinent. For these reasons we also have confidence in our retrieved cooling trends over for example daytime continental Central/Eastern Africa, which are different from the other four day/night datasets.

The spatial correlations between AIRS_RT retrieved rates and the various datasets is shown in Table 2 while the cosine weighted skin temperature trends are shown in Table 3. By adding in the uncertainty in the trends for any of the individual models or datasets,




Figure 5. Surface temperature trends $dSKT/dt$ averaged over day and night for AIRS_RT, and from separately fitting the monthly data in ERA5, MERRA2, AIRS L3, CLIMCAPS L3 and GISS. The horizontal and vertical axis are longitude and latitude. Colorbar units are in $K\ yr^{-1}$.

and then doing the cosine weighting, we estimate uncertainties of about $\pm 0.015\ K\ yr^{-1}$ for “ALL”; the uncertainties for “OCEAN” are typically about 2/3 of that value, and for “LAND” are about 4/3 of that value. We emphasize here that we use center point **NWP reanalysis** and L3 model data when computing their trends for any grid box, while the AIRS_RT uses the hottest 10% of “clear” **data observations**; (Strow & DeSouza-Machado, 2020) showed that the tropical retrieved surface temperature trends and anomalies over ocean correlated very well with those from the ERA-I Sea Surface Temperature dataset.

ERA5	MERRA2	AIRSL3	CLIMCAPSL3	GISS
0.72	0.59	0.80	0.89	0.77

Table 2. Correlations of average (nighttime, daytime) retrieved skin temperature trends from AIRS_RT, versus trends from models/products

A notable outlier in this group is the MERRA2 trends, especially over land and the Southern Ocean which are noticeably negative (blue) compared to the other datasets; the agreement with tropical and mid-latitude oceans is much better. As noted earlier, the MERRA2 monthly trends come from a combination day/night dataset that was downloaded, which as seen in Figure 5 consists of trends that are both positive and negative, combining to get a closer-to-zero global weighted trend. In addition MERRA2 is the only one of the six that (a) does not have the extreme $+0.15\ K\ yr^{-1}$ warming in the northern polar region and (b) shows **a lot of substantially more** cooling in the Central African

SKT trend K yr ⁻¹	AIRS_RT	AIRS	CLIMCAPS	ERA5	MERRA2	GISS
ALL	0.020	0.017	0.021	0.023	0.011	0.021
TROPICS	0.011	0.011	0.012	0.016	0.010	0.015
MIDLATS	0.029	0.020	0.028	0.026	0.020	0.026
POLAR	0.032	0.028	0.033	0.041	-0.005	0.028
OCEAN	0.019	0.011	0.019	0.017	0.012	0.017
LAND	0.022	0.030	0.024	0.038	0.010	0.030

Table 3. Cosine weighted skin temperature trends (in K yr⁻¹); uncertainties are on the order of ± 0.015 K yr⁻¹ as explained in the text.

area. Using ERA5 monthly data, we devised a test similar to the one mentioned in Section 4 to determine if the differences between MERRA2 and ERA5 surface temperature trends could be due to the temporal sampling (once for MERRA2 versus eight times for ERA5). For each month we matched the eight ERA5 timesteps available per month to the tile centers and then averaged the surface temperatures per month; the ensuing geophysical timeseries was then trended. The day/night ERA5 average of Figure 5 was compared to these trends; of note are (a) we did not see the cooling in Africa and near the Antarctic that is seen in MERRA2 and (b) the main differences between the 1.30 am/1.30 pm average in the bottom middle (ERA5) panel were over land (all 5 continents); the histograms of the differences showed the peak was typically close to 0 K yr⁻¹, but the widths over land were about ± 0.02 K yr⁻¹ or less (compared to ± 0.005 K yr⁻¹ over ocean). Both AIRS L3 and MERRA2 show cooling in the Southern Ocean; we note that although MERRA2 is the *a-priori* for CLIMCAPS L3, their trends are different than those from MERRA2; in fact AIRS_RT shows the closest correlation to the observational CLIMCAPS L3 trends. The AIRS L3 trends in the Southern Ocean region could arise because of problems identifying ice during the L2 retrieval (private communication : Evan Manning (JPL) and John Blaisdell (NASA GSFC)) though the MERRA2 trends also show significant cooling in that region, where few surface observations from buoys poleward of 60° exist to help resolve these differences (see for example Figure 10 in (Haiden et al., 2018)).

Figure 6 shows the zonally averaged total (land+ocean) and ocean only surface temperature trends. The equator to midlatitude ocean trends are almost linear for all datasets, with the slope for the northern hemisphere being about double that of the southern hemisphere (roughly 0.001 K yr⁻¹ per deg latitude). Again focusing on the right hand plot, the AIRS L3 trends are negative in the Southern Ocean regions, compared to the other 3 datasets, due to the cooling trends around the Antarctic continent shown earlier, but then agrees with most of the other datasets over the Antarctic; the MERRA2 trends significantly differ between -90 S and -50 S. MERRA2 and ERA5 also show slightly smaller warming trends in the Northern Polar, compared to the three ~~AIRS-based~~ AIRS observation-based datasets.

We point out that the trends seen in Figure 5 vary noticeably at more local, regional levels and furthermore this spatial variation can differ between daytime and nighttime, evident in Figure A1 of ~~Appendix~~ Appendix A, and that the observational sets (AIRS_RT, CLIMCAPS L3 and AIRS L3) had larger differences than ERA5. Discussing the possible causes is outside the scope of the paper.




Figure 6. Zonally averaged surface temperature trends for (left) sum of ocean and land point and (right) ocean only. The vertical units are K yr^{-1} while the horizontal axis are degrees latitude.




Figure 7. dWV_{frac}/dt (left) without and (right) with MLS *a-priori* in the upper atmosphere. The vertical axis are pressure (in mb), the horizontal axis are latitude (in degrees) while the colorbar is in yr^{-1} (fractional water vapor has no units).

6.2 Addition of Microwave Limb Sounder Water Vapor A-priori

The Microwave Limb Sounder (MLS), on board NASA's Aura platform, flies about 15 minutes behind AIRS on the same orbit. It is designed for sounding of the atmosphere above 300 mb. We computed water vapor trends from the L3 data produced for that instrument (above 300 mb) and used them as an *a-priori* for the AIRS_RT retrieval.

Figure 7 shows the retrieved fractional water vapor trends when the *a-priori* trend in the upper atmosphere in the left and right panels were zero, or used MLS trends, respectively. One sees that the additional information brought in by the instrument sensitive to upper troposphere humidity, significantly changes the water vapor sounding especially in the polar region by moving towards the MERRA2 and ERA5 fractional water vapor trends seen in Figure 10. We note that the other related results shown in this paper also use the MLS *a-priori*.




Figure 8. Zonally averaged column water vapor trends for AIRS_RT, AIRS L3, CLIMCAPS L3, ERA5 and MERRA2. Vertical units are in mmw yr⁻¹ while the horizontal axis are in degrees latitude.

6.3 Column water vapor trends

Column water is dominated by water vapor amounts close to the surface and the column vapor trends thus provide an assessment of the water vapor retrieval quality in the lower atmosphere. The water vapor information in the lowest layers is best retrieved using the weak water lines in thermal infrared region. As noted earlier this part of the retrieval is significantly complicated by the simultaneous presence of nonzero surface temperature, air temperature and water vapor jacobians in this spectral region, meaning the AIRS instrument has much reduced sensitivity to the water vapor amounts in these lowest layers. In addition the changing concentration of very minor gases such as CFC-11 and CFC-12 (Strow & DeSouza-Machado, 2020) are quite evident in the spectral trends, further complicating the water vapor trend retrieval for the lowest layers.

Figure 8 shows the zonally averaged column water vapor trends; not shown are the error bars which are on the order of ± 0.005 mm/year. AIRS_RT is from our retrievals while the rest are directly from the NWP-reanalysis or L3 model-fields. Close examination shows the CLIMCAPS L3 column water trend is nearly identical to the MERRA2 trend, as is also seen in lower atmosphere water vapor trends shown later in Figure 10. Conversely the column water vapor trends for AIRS L3 are negative in the lower troposphere in the midlatitudes and tropics, which is not to be expected given that the surface temperature trends are positive. AIRS_RT nominally agrees with ERA5 and MERRA2 in the tropics and midlatitudes, but is smaller than either in the northern polar regions. A reduced rate for AIRS_RT is additionally seen in the 0-50 N latitudes, where there is a larger fraction of land (for which we do not use the assumption of constant relative humidity) compared to the Southern Hemisphere. Screening out the tiles over land slightly improves the agreement between reanalysis (ERA5, MERRA2) vs AIRS_RT column wa-

ter trends. Examination of the spectral trends in the window region does not shed any more insight into the differences, as the observation spectral trends and ~~NWP-reanalysis~~ reconstructed trends are very similar and we are fitting the observed trends. The magnitudes and patterns look similar to the 2005-2021 column water trends shown in (Borger et al., 2022), which were derived using observations from the Ozone Monitoring Instrument (OMI). We point out their 16 year zonally averaged trends look similar to the 20 year ERA5 zonally averaged column water trends between -60°S and -10°S , but become almost a factor of 2 larger between -10°S and $+40^{\circ}\text{N}$; the zonally averaged OMI 16 year trends are negative in the polar regions. The column water trends are summarized in Table 4.

DATASET mm yr ⁻¹	OMI 16 years	AIRS_RT 20 years	ERA5 20 years	MERRA2 20 years	AIRS L3 20 years	CLIMCAPS L3 20 years
GLOBAL (cosine average)	0.051	0.021	0.035	0.036	-0.009	0.038
TROPICAL	0.083	0.028	0.047	0.042	-0.015	0.045

Table 4. Column water trends based on OMI ~~data~~ observations (16 years) and AIRS_RT, ERA5 and MERRA2 (20 years). The units are in mm yr⁻¹; the uncertainties are on the order of 0.1 mm yr⁻¹ for OMI and AIRS_RT, and half that for ERA5 and MERRA2, and AIRS L3 and CLIMCAPS L3.

D/N differences (not shown) for AIRS_RT were on the order of ± 0.005 mm yr⁻¹ (with daytime trends being smaller over land), for AIRS L3 were on the order of ± 0.01 mm yr⁻¹ or more (with larger values happening over the daytime tropical oceans), while that for ERA5 and CLIMCAPS L3 were typically on the order of ± 0.03 mm yr⁻¹ or less.

6.4 Zonal atmospheric temperature and water vapor trends

Figure 9 shows the zonally averaged atmospheric temperature trends from five of the datasets in Figures 5,8 above. In the troposphere the AIRS_RT retrievals show the same general features as the trends from ERA5, though they begin to diverge in the stratosphere and especially above that. In particular AIRS_RT does not show warming in the Southern Polar stratosphere; we have separately looked into seasonal trends and noted that our retrieved September/October/November temperature trends in the upper atmospheric Southern Polar regions are on the order of -0.12K yr^{-1} , possibly leading to an overall no net heating/cooling for the annual trends. We highlight that our results are smoother than those of the other datasets, while the other sets have noticeable discontinuities that may not be physical under the thermodynamics or fluid dynamics frameworks. In addition we point out that both our results and AIRS v7 L3 show a hint of cooling over the tropical surfaces. Note that CLIMCAPS is initialized by MERRA2, and their temperature trends are quite similar. AIRS v7 looks similar to AIRS_RT except in the tropics where it almost has cooling in the lower troposphere and much more warming in the lower stratosphere. The correlations between AIRS_RT and the [AIRS L3, CLIMCAPS L3, MERRA2, ERA5] temperature trends of Figure 9 are [0.74,0.65,0.74,0.72] respectively.




Figure 9. Zonally averaged dT/dt shown in 5 panels. Horizontal axis is in degrees latitude while vertical axis is pressure (mb). The y -limits are between 10 to 1000 mb, on a logarithmic scale. The colorbar is units of in $K\ yr^{-1}$.




Figure 10. Zonally averaged dWV_{frac}/dt shown in 5 panels. Horizontal axis is latitude while vertical axis is pressure. The y -limits are between 100 to 1000 mb, on a linear scale. [The colorbar units are in \$\text{yr}^{-1}\$, as fractional water vapor is dimensionless.](#)




Figure 11. The 400 mb fractional water vapor trends for (left) AIRS_RT and (right) ERA5 show general agreement except in the Southern Polar Regions. The colorbar units are in yr^{-1} , as fractional water vapor is dimensionless.

Figure 10 shows the zonally averaged atmospheric fractional water vapor trends ($d/dt \text{ WV}(z,t)/\langle \text{WV}(z,t) \rangle$). The five panels are markedly different from one another. The AIRS_RT trends resemble those of ERA5 in the tropical troposphere, though we do not have drying in the lower tropical layers. Conversely, the observed trends in the Southern Polar (AIRS L3, CLIMCAPS L3 and AIRS_RT) show drying rather than wetting, though AIRS_RT is less than that of CLIMCAPS/MERRA2. AIRS_RT is an outlier in the upper polar atmosphere trends, as both the signals and the jacobians are close to zero. Of some concern is a little bit of drying in the northern polar region, where there are low H_2O amounts leading to small jacobians. CLIMCAPS v2 looks quite similar to the MERRA2 trends. AIRSv7 shows substantial drying in the lower troposphere, and considerable wetting in the upper troposphere, compared to any of the other datasets. Spectral closure studies (using the AIRS v7 H_2O trend \times the H_2O jacobians derived above from ERA5 average profiles) are not shown here, but differ noticeably from the CCR trends from AIRS v7 in the $1300\text{-}1600 \text{ cm}^{-1}$ region, indicating there are inadequacies in the AIRS V7 water vapor retrievals. The correlations between AIRS_RT and the [AIRS L3, CLIMCAPS L3, MERRA2, ERA5] fractional water vapor trends of Figure 10 (limited to 100 mb, 1000 mb) are [0.65, 0.24, 0.36, 0.58] respectively.

Figure 11 shows the 400 mb fractional water vapor trends, with the left panel being the AIRS_RT trends while the right panel is the ERA5 trends. Note that there is general agreement except in the Southern Polar region, as also seen later in Figure 10 in the other two observational L3 datasets (AIRS v3 and CLIMCAPS). This could be related to work by (Boisvert et al., 2019) who showed decreasing evaporation from the Southern Ocean in the 2003-2016 period due to increasing ice cover.

7 Uncertainty

The uncertainties for the AIRS v7 geophysical products are impacted by radiance noise amplification due to cloud clearing (Susskind et al., 2003) and the neural net first guess, while state vector errors are estimated based on regressions. CLIMCAPS L2 geophysical products are similarly impacted by cloud clearing noise in the radiances, but these are fully propagated together with geophysical error estimates from the MERRA2 first guess, through the retrieval algorithm which uses Optimal Estimation (Smith & Barnett, 2020). No estimate of uncertainties are available for the monthly L3 products.

The uncertainties for the AIRS_RT trends is much more straightforward : the spectral uncertainties shown in Figure B2 are used together with the state vector covariance matrices to generate the uncertainty matrix using the relevant equations of Optimal Estimation Rodgers (2000); we use the diagonal elements for the final uncertainties. Panels (A) and (C) of Figure 12 shows the zonally averaged (D/N) uncertainties as a function of pressure and latitude. Inspection of the radiance trends uncertainties shown in the center panel of Figure B2 shows the upper atmosphere temperature sounding region (650-700 cm^{-1}) has much larger uncertainty in the polar regions. The instrument and spectroscopy characteristics, coupled with these observational uncertainties, are such that for temperature the smallest errors are in the tropics while the largest errors are in polar upper atmosphere, which are the regions below 100 mb where the ERA5 trends differ most from AIRS_RT trends. Similarly for water vapor the larger errors are in the lower atmosphere and above about 300 mb; the constant RH assumption and MLS *a-priori* help alleviate the errors in the retrieved trends. [We point out earlier work on studying upper tropospheric/lower stratospheric humidity over tropical cyclones also used MLS climatology together with AIRS observations](#) (Feng & Huang, 2021).

The *Z-test* confirmed this picture, as seen in panels (B) and (D) of Figure 12, which show the temperature and fractional water vapor trends, together with black dots marking the (latitude, altitude) points where the trends are larger than the uncertainty in the trends, at the 5% significance level. This happens in panel (B) for the temperature trends in most of the tropical/mid-latitude free troposphere (and stratosphere) but not at the southern polar stratosphere; and in panel (D) for fractional water vapor trends in the 200-600 mb range, from the Southern Polar region to about +60 N latitude, and some spots in the Northern Polar.

8 Discussion

In general for surface temperature trends, the disagreements between the six sets shown in Figure 5 are over the polar regions and over land (especially over the Amazon and Central Africa) and are smallest over tropical and mid-latitude oceans, indicating the best agreements, except for slightly larger differences off the western coast of the Americas and Africa (which have a prevalence of MBL clouds). The atmospheric temperature trends in general agreed except for the upper atmosphere polar regions and in the high altitudes (less than about 200 mb). Similarly fractional water vapor trends differed most in the upper atmosphere (200 mb and above) and in the tropical/mid-latitude 600-800 mb region. A quick glance at Figure 10 shows the former is due to lower sensitivity to upper atmosphere water vapor, leading the AIRS_RT retrievals to have low values while the AIRS L2 retrieval is initialized by a neural net; conversely the latter is due to the AIRS L3 retrieval being negative while the rest were mainly positive. Similarly the AIRS_RT retrieval differs above the Antarctic continent.

In general the observed surface temperature trends from the AIRS_RT retrievals agree with the ERA5 and MERRA2 trends, as well as the NASA GISS trends, except in the Southern Antarctic. That is a region where there are few surface observations; for retrievals there are competing effects of using ice vs ocean surface emissivity. Overall, the AIRS_RT retrieved surface temperature trends are typically in between ERA5 and MERRA2 for land + ocean in all regimes (tropical, midlatitude and polar), though slightly larger overall for ocean than the two reanalysis datasets; in general they are closer to the ERA5 trends than the MERRA2 trends.

(Strow et al., 2021) demonstrated that the long- and medium- wave channels of the AIRS instrument are radiometrically stable to better than $0.002\text{-}0.003 \text{ K yr}^{-1}$, which is much smaller than the surface and tropospheric temperature trends in the reanalysis mod-




Figure 12. Zonally averaged D/N plots of (A) temperature uncertainties in K yr^{-1} and (B) temperature trends in K yr^{-1} together with null hypothesis. (C) and (D) are the same except for fractional water vapor uncertainty and trends in $1/\text{year yr}^{-1}$. Horizontal axis are in degrees latitude while vertical axis are pressure (mb) - logarithmic for temperature and linear for water vapor. See text for more detailed explanation.

els, AIRS L3 data and our retrieved trends. A separate analysis of spectral trend uncertainties after 05,10,15,20 years (not shown here) show that these uncertainties have been steadily decreasing and are now approaching this number, as can be seen in the bottom left panel of Figure 3. Furthermore, though we cannot guarantee only cloud free scenes in our chosen Q0.90 observational dataset used in this paper, the high correlations between other dataset surface trends compared to ours, is a good indication that our results come from mostly cloud-free scenes, or scenes whose clouds have negligible impact on our results.

The observed zonal temperature trends agree with those from the models and the AIRS L3 products, except in the polar regions. Again this could be an issue of using slightly incorrect surface emissivity for the AIRS_RT retrievals. In addition we point out that since there is very little water vapor, the temperature jacobians near the surface are quite small in magnitude (compared to more humid atmospheres) and so it is difficult to separate out the effects of surface temperature trends versus lower atmosphere temperature and H₂O trends. The quantile construction used in this paper means that for example tiles straddling the subcontinent of India and the ocean will preferentially pick the land surface observations for daytime, which could lead to misleading trends on these coastal tiles. It is possible to subdivide the $3^\circ \times 5^\circ$ tiles into for example $1^\circ \times 1^\circ$ grids and do the analysis, but the number of observations per small grid cell would drop, leading to more noise in the retrieved trend.

In general the AIRS_RT retrieved ~~absolute~~ column water trends are slightly smaller than ERA5 in the Southern Hemisphere but noticeably smaller in the Northern Tropics to midlatitudes. We have mentioned difficulties we have retrieving H₂O close to the surface and in the upper atmosphere, due to the known sensitivity of infrared sounders whose water vapor averaging kernels peak in the 300-600 mb range, and we have pointed out examination of the spectral residuals in the window region shows we are fitting the signal. The MERRA2 and CLIMCAPS column water vapor trends are quite similar, while the AIRSv3 L3 trends are noticeably different, being negative almost everywhere. If we start with zero *a-priori* for water vapor at the surface, we can fit the spectral trends but the retrieved water vapor trends in the lower layers which dominate column water amounts, can leads to column water trends that are easily double or more than the results for the other datasets.

Given the complex numerical algorithms used in both the reanalysis models and the AIRS L3 retrievals as well as those in the AIRS_RT trends, it is difficult to offer precise explanations for any of the trends shown above. Our results are relatively robust to changes in the covariance or Tikonov parameter settings. For instance changing them by factors of two would keep the trends about the same, though of course the uncertainties would change. There are however a few general points that can be made. The first is that since infrared instruments are sensitive to the 300-800 mb region and lose sensitivity outside this, the retrievals from AIRS_RT and AIRS L3 have difficulties with water vapor in the lower (Planetary Boundary Layer) and upper troposphere/lower stratosphere. One way to mitigate this is to use trended observed data from external sources in the *a-priori*, while keeping the *a-priori* trends for all other parameters as 0. For example we have shown we can use the MLS ~~data~~ observations above 300 mb without significantly degrading the AIRS_RT retrieval in the middle and lower atmosphere; conversely the CLIMCAPS retrievals are initialized by MERRA2 and while they can pull out weather signals, their L3 trends are still quite closely tied to the MERRA2 trends. The tropical and mid-latitude ocean surface temperature trends from the numerical models that assimilate observed data, L3 products and AIRS_RT are very similar; however they start to show differences where there are few *in-situ* ~~data~~ observations combined with problems with ice identification (surface emissivity)/cold temperatures which exacerbate the drifting AIRS detector problems (Strow et al., 2021), such as the Arctic and Southern Ocean.

9 Conclusions

We have ~~described a novel method~~ designed a novel retrieval method, specifically to obtain global thermodynamic atmospheric climate trends, ~~starting from~~. It uses longterm stable, high spectral resolution infrared allsky hyperspectral observations which are ~~then~~ first subset for “nominally clear” scenes. ~~Our retrieved~~ The geophysical trends are derived using ~~trends from observed trends from the~~ well characterized (radiometrically stable) radiances and from zero *a-priori* (except for a constant relative humidity assumption). This makes them much more direct and traceable than trends from traditional L2 retrieval algorithms, which use complicated *a-priori* information. We also ~~did performed~~ “radiative closure” tests by running the monthly ~~NWP reanalysis~~ or L3 ~~fields data~~ through a radiative transfer model to compare the spectral trends so obtained against the observed spectral trends, ~~which showed the most disagreement in~~. The most noticeable disagreement in spectral trend radiance space was in the water vapor free troposphere sounding regions.

The temperature and water vapor trends retrieved from the “nominally clear” radiance trends resemble those computed from monthly ERA5 and MERRA2 reanalysis. The radiative spectral closure helps identify the cause of differences in the geophysical trends, rather than solely attributing them to deficiencies (eg the well known reduced sensitivity to water vapor near the boundary layer and above 200 mb) with our retrieval. For example the AIRS_RT temperature trends are quite similar to the reanalysis (MERRA2/ERA5) trends, while the water vapor (and/or Relative Humidity) trends are quite different, especially in the lower troposphere and upper troposphere, which is clearly manifest as differences in the spectral trends in the water vapor sounding region.

The 20 years of AIRS observations were binned into nominal 3×5 degree grid boxes covering the planet, with a time step of 16 days, from which anomalies and trends were obtained. To alleviate the reduced sensitivity of hyperspectral sounders to water vapor in the lower atmosphere we used an assumption of 0.01 increase in relative humidity to initialize the *a-priori* lower atmosphere fractional water vapor rates, while we similarly used Microwave Limb Sounder trends as an *a-priori* to address the high altitude water vapor deficiencies caused by lower sensitivity to upper atmosphere water vapor. New or updated time dependent surface emissivity databases may become available in the future, enabling us to include those effects into Equation 4. Problems in the polar regions and Planetary Boundary Layer water vapor retrievals will be harder to overcome since there is very little sensitivity to water vapor in these regions, together with fewer observations to compare against, though more work is planned to address both of these.

In this paper we used the 90th quantile (Q0.90) nominally “hottest” observed BT1231 ~~data~~ to form a time series over which to obtain radiance trends, after establishing that the spectral trends from this quantile differed by less than about $\pm 0.0015 \text{ K yr}^{-1}$ from the 50th (or average) quantile. In the future we plan to base the ~~data~~ subset selection on MODIS cloud products (obtained at 1 km resolution compared to the AIRS 15 km resolution). In any case the AIRS L1C Q0.90 spectral trends used for the AIRS_RT results are very comparable to trends from quality assured binned AIRS CCR data (Manning, 2022). The quantile method allows us to select which ~~data~~ observations to use in the trends : we have explored doing the trend retrievals using the cloud fields contained in ERA5, together with the TwoSlab cloud algorithm (De Souza-Machado et al., 2018) to compute jacobians when clouds are present, together with trends from the Q0.50 observational dataset described above. The retrieved geophysical trends resemble those described above in the mid to upper atmosphere, and differ in the lower atmosphere, but more work is needed on this and is not discussed further. Longwave clear sky flux trends (both outgoing top-of-atmosphere and incoming bottom-of-atmosphere) and climate feedbacks will be discussed in a separate paper.

While the Aqua platform is scheduled to be terminated within the next few years, copies of near identical CrIS instruments are already in orbit, and more will be launched over the next few years, till at least 2040. The Climate Hyperspectral Infrared Radiance Product (CHIRP) (Strow et al., 2021) will seamlessly combine the AIRS ~~data~~ observations between 2002-2015 to CrIS ~~data~~ observations from 2015-2040 to obtain a 40 year observational radiance record over which to study climate. This availability means that AIRS_RT and future AIRS/CrIS versions, is well positioned to enable climate analysis of geophysical trends for years to come.

Appendix A Day versus Night surface temperature trend differences

Figure A1 shows the (top) daytime and (middle) nighttime surface temperature trends; from left to right the datasets are (observational) AIRS_RT, AIRS L3, CLIMCAPS L3 and (reanalysis) ERA5. In general the AIRS ~~observational datasets~~ observations show enhanced daytime cooling over the Indian subcontinent and Central Africa, compared to the ERA5 model; they also show daytime warming trends over continental Europe and central Asia and the Amazon are larger than during the nighttime. With the large ocean heat capacity and smaller land heat capacity, the land is expected to show more of a diurnal cycle than ocean. ERA5 sees warming over Eastern/Central Africa during daytime while the ~~observational datasets~~ see observations show cooling. Similarly the three ~~observational datasets~~ see observations show more daytime cooling over the Indian sub-continent and south eastern Australia than does ERA5; we omit more detailed analysis in this paper. During the nighttime, the AIRS L3 product has cooling over C. Africa and parts of the Amazon. The day-night differences are seen in the bottom row of the same figure. Note the colorbar is the same for all three rows. The differences are close to zero over the ocean. AIRS_RT and CLIMCAPS L3 see more daytime cooling over E. Africa and the Indian subcontinent. Overall the magnitude of the day - night differences for the observations are larger for the AIRS ~~observational datasets~~ observations than for ERA5. ERA5 also sees negative differences over Central Asia compared to the AIRS ~~observational datasets~~ observations, which see positive differences (higher surface temperature trends during the daytime).

The atmospheric temperature and fractional water vapor day-night differences are quite small (compared to the average values) and not shown here; AIRS L3 shows noticeable more wetting of the 600-800 mb region during daytime versus nighttime, compared to the other three.

Appendix B Spectral closure : comparisons between observed and simulated spectral trends

The main body of the manuscript details comparisons of climate geophysical trends using a purpose designed algorithm to analyze radiance observation trends, versus those from reanalysis and monthly L3 fields. In this Appendix we present the comparisons in radiance spectral trend space, by using the spectral closure method to assess monthly thermodynamic output from reanalysis and/or L3 products (see for example (X. Huang et al., 2023)). This is accomplished by geolocating the entire 20 year monthly reanalysis and L3 surface temperature, air temperature, water vapor and ozone fields for all 72×64 tiles. We also include realistic column linearly-increasing-with time mixing ratios for CO_2 , CH_4 and N_2O as well as land or ocean surface emissivity co-located to tile centers together with view angles of about 22° , which is the average view angle of the tiled observations. The model fields are then converted to spectral radiances by running through the SARTAfast model (Strow, Hannon, DeSouza-Machado, et al., 2003). Finally, spectral radiance trends are computed from the time series of (clear sky) spectral radiances using Equation 2.




Figure A1. Top two rows : The (top) day and (middle) night surface temperature trends for AIRS_RT, AIRS L3, CLIMCAPS L3 and ERA5. Third row (bottom) is the D-N difference. Colorbar units are in $K\ yr^{-1}$.

Here we select two examples to illustrate differences in the five datasets we use in this paper. Firstly we study spectral trends in the water vapor sounding region. Water vapor is highly variable in space and time, meaning water vapor retrievals using hyperspectral sounders radiances differ most from Numerical Weather Prediction (NWP) forecasts. In particular the typical ± 90 minute difference between observation and forecast means sounders provide most accurate water vapor information, when considered locally and at a particular time. However this will not affect the water vapor trends we show in this paper since atmospheric water vapor timescale is on the order of about a week to ten days (van der Ent & Tuinenburg, 2017), and we are also considering data points averaged over 16 or more days. Figure B1 show the globally averaged brightness temperature trends (in $K\ yr^{-1}$) in the $1350 - 1650\ cm^{-1}$ water vapor sounding region. The blue curve shows the trends from the AIRS observations used in this paper, while spectral trends constructed from the AIRS L3/ CLIMCAPS L3 retrievals are in red/yellow and the ERA5 model fields are in purple. The AIRS observations and ERA5 constructed spectral trends are positive in this region, while the AIRS L3 and CLIMCAPS L3 trends are obviously different, being negative in this water vapor sounding region. The subtle differences in these spectral trends arise from differences in the geophysical trends between observations and the models themselves, and were addressed in Sections 6.3 and 6.4 of the text.

Data Availability Statement

The observation data-




Figure B1. Globally averaged spectral trends (in K yr^{-1}) for the $6.7 \mu\text{m}$ ($1400\text{--}1650 \text{ cm}^{-1}$) free troposphere water vapor sounding region, as a function of wavenumber (cm^{-1}). AIRS L1C observations (blue) are compared to spectral closure from the standard monthly AIRS L3 retrievals (red) and CLIMCAPS L3 (yellow) and from monthly ERA5 simulations (purple). The reconstructed AIRS_RT trends very closely match the AIRS L1C observations and are not shown here.




Figure B2. 20 year zonally averaged spectral brightness temperature trends (colorbars in K yr^{-1}) for nighttime (left) AIRS Q0.90 observations and (right) clear sky simulations using ERA5 monthly model fields. The center panel shows the AIRS Q0.90 spectral uncertainties (colorbar also in K yr^{-1}). Realistic linear trends of CO_2 , CH_4 and N_2O were included in the ERA5 simulations, while the O_3 trends in ERA5 are from the reanalysis itself. Horizontal axis are in wavenumbers (cm^{-1}) while vertical axis are in degrees latitude.

1350

1351 Second, we focus on comparing zonally averaged spectral trends between AIRS observations
 1352 and ERA5 simulations. Figure B2 shows the AIRS observed Q0.90 (nominally clear) descending
 1353 (night) zonally averaged results in K yr^{-1} in the left panel, and the zonally averaged simulated
 1354 clearsky (without clouds) spectral trends (also in K yr^{-1}) from monthly ERA5 fields in
 1355 the right panel. The center panel shows the spectral trend uncertainties from the observations,
 1356 also in K yr^{-1} . Earlier sections, including Section 6.4 compared the geophysical trends
 1357 between retrieved from AIRS observation and reanalysis/L3 data fields. The similarities/
 1358 differences in geophysical trends between observations and models/operational data can
 1359 be partially understood from the similarities/differences in the spectral trends. For example,
 1360 the H_2O sounding region ($1350\text{--}1600\text{ cm}^{-1}$) of the left and right panels of Figure B2 shows
 1361 roughly similar (positive) spectral trends in the tropics and mid-latitudes; there are some
 1362 slight differences in the high altitude channels ($1450\text{--}1550\text{ cm}^{-1}$ region). The main body
 1363 of text demonstrated how these differences translate to subtle differences in the geophysical
 1364 trends. Observations and simulations both have positive dBT/dt in the $800\text{--}960, 1150\text{--}1250$
 1365 cm^{-1} region, indicating surface warming; however the ERA5 simulation show more warming
 1366 in the southern polar regions than do the AIRS observations. Note the mean warming
 1367 in the tropics for both observations and ERA5 simulations is less than that in the mid-latitudes,
 1368 and the polar regions show the largest overall change in brightness temperature in the
 1369 window region. Large differences are seen in the $10\text{ }\mu\text{m}$ (1000 cm^{-1}) O_3 sounding region,
 1370 which are not surprising since ozone assimilation is not a primary goal of ECMWF assimilation;
 1371 here we do not address these as we focus on the changes to the moist thermodynamic
 1372 state. The window region trends computed using the ERA5 model are more positive in
 1373 the Southern Polar region. Conversely the $640\text{--}700\text{ cm}^{-1}$ spectral region is positive, especially
 1374 in the tropics; however the observations show a net cooling trend away from the tropics,
 1375 compared to the ERA simulations. This demonstrates the importance of the model \rightarrow
 1376 spectral trend comparisons, given the accuracy of the AIRS observations.

1377 Data Availability Statement

1378 The observations used in this paper (AIRS L1C radiances), as well as the AIRS L3,
 1379 CLIMCAPS L3, MERRA-2 and Microwave Limb Sounder monthly data products, are
 1380 freely available to the public on the NASA Goddard Space Flight Center Earth Sciences
 1381 (GES) Data and Information Services Center (DISC) servers <https://disc.gsfc.nasa.gov/>.
 1382 Monthly ERA5 is freely available through (single levels) [https://cds.climate.copernicus.eu/](https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels-monthly-means?tab=overview)
 1383 [datasets/reanalysis-era5-single-levels-monthly-means?tab=overview](https://cds.climate.copernicus.eu/datasets/reanalysis-era5-single-levels-monthly-means?tab=overview) and (pressure lev-
 1384 els) [https://cds.climate.copernicus.eu/datasets/reanalysis-era5-pressure-levels-monthly-](https://cds.climate.copernicus.eu/datasets/reanalysis-era5-pressure-levels-monthly-means?tab=overview)
 1385 [means?tab=overview](https://cds.climate.copernicus.eu/datasets/reanalysis-era5-pressure-levels-monthly-means?tab=overview). GISTEMP monthly model output are also freely available from
 1386 <https://data.giss.nasa.gov/gistemp/>. The Matlab based source code used for the anal-
 1387 ysis is freely available on https://github.com/sergio66/oem_climate_code, while the F90
 1388 kCARTA (De Souza-Machado et al., 2018) line-by-line code used to make the jacobians
 1389 is freely available on https://github.com/sergio66/kcarta_gen.

1390 Acknowledgments

1391 We gratefully acknowledge the help of H. Motteler, who designed, implemented and
 1392 ran the AIRS tiling code. C. Hepplewhite helped test the outputs of the tiling code, S.
 1393 Buczkowski worked on bringing down almost all the data used in this study, and made
 1394 the uniform/clear [observational](#) datasets. Chris Barnett has given helpful comments and
 1395 general advice regarding retrievals and comparisons to the AIRSv7 and CLIMCAPS datasets.
 1396 [Jing Feng provided valuable feedback regarding the manuscript](#). The hardware used is
 1397 part of the UMBC High Performance Computing Facility (HPCF). The facility is sup-
 1398 ported by the U.S. National Science Foundation through the MRI program (grant nos.
 1399 CNS-0821258, CNS-1228778, OAC-1726023, and CNS-1920079) and the SCREMS pro-

1400 gram (grant no. DMS-0821311), with additional substantial support from the Univer-
1401 sity of Maryland, Baltimore County (UMBC). See hpcf.umbc.edu for more information
1402 on HPCF and the projects using its resources.

References

- 2023, G. T. (2005). *GISS Surface Temperature Analysis (GISTEMP), version 4* (Tech. Rep.). data accessed 2023/11/30 at <https://data.giss.nasa.gov/gistemp/>: NASA Goddard Institute for Space Studies.
- Allan, R., Willett, K. M., John, V., & Trent, T. (2022). Global Changes in Water Vapor 1979–2020. *J. Geophys. Res.*, *127*, e2022JD036728. <https://doi.org/10.1029/2022JD036728>.
- Anderson, T., Hawkins, E., & Jones, P. (2016). CO₂, the greenhouse effect and global warming: from the pioneering work of Arrhenius and Callendar to today's Earth System Models. *Endeavour*, *40*, 178–187; <https://doi.org/10.1016/j.endeavour.2016.07.002>.
- Andersson, E., Holm, E., Bauer, P., Bejaars, A., Kelly, G., McNally, A., ... Tompkins, A. (2007). Analysis and forecast impact of the main humidity observing systems. *Quart. J. Roy. Meteorol. Soc.*, *133*, 1473–1485; <https://doi.org/10.1002/qj.112>.
- Arrhenius, S. (1896). On the influence of carbonic acid in the air upon the temperature of the ground. *London, Edinburgh and Dublin Phil. Mag. and J. of Sci.*, *5*, 237–276; <https://doi.org/10.1080/14786449608620846>.
- Boisvert, L., Vihma, T., & Shie, C.-L. (2019). Evaporation From the Southern Ocean Estimated on the Basis of AIRS Satellite Data. *JGR*, *125*, <https://doi.org/10.1029/2019JD030845>.
- Borbas, E., Hulley, G., Feltz, M., Knuteson, R., & Hook, S. (2018). The Combined ASTER MODIS Emissivity over Land (CAMEL) Part 1: Methodology and High Spectral Resolution Application. *Remote Sens.*, *10*, 643; <https://doi.org/10.3390/rs10040643>.
- Borger, C., Beirle, S., & Wagner, T. (2022). : Analysis of global trends of total column water vapour from multiple years of OMI observations. *Atmos. Chem. Phys.*, *22*, 10603–10621, <https://doi.org/10.5194/acp-22-10603-2022>.
- Brindley, H., & Bantges, R. (2016). The Spectral Signature of Recent Climate Change. *Curr. Clim. CHange. Reports*, *2*, 112–116.
- Brindley, H., Bantges, R., Russell, J., Murray, J., Dancel, C., Belotti, C., & Harries, J. (2015). The Spectral Signature of Earth's Climate Variability over 5 years from IASI. *J. Clim.*, *28*, 1649–1160; DOI: 10.1175/JCLI-D-14-00431.1.
- Bromwich, D., Ensign, A., Wang, S.-H., & Zou, X. (2024). Major Artifacts in ERA5 2-m Air Temperature Trends Over Antarctica Prior to and During the Modern Satellite Era. *GRL*, *51*, <https://doi.org/10.1029/2024GL111907>.
- Cai, M., & Kalnay, E. (2004). Can Reanalysis Have Anthropogenic Climate Trends without Model Forcing? *J. Clim.*, *18*, 1844–1849.
- Carn, S., Strow, L., DeSouza-Machado, S., Edmonds, Y., & Hannon, S. (2005). Quantifying tropospheric volcanic eruptions with AIRS : the 2002 eruption of Mt. Etna (Italy). *Geophys. Res. Lett.*, *32*, L02301; doi:10.1029/2004GL021034.
- Chedin, A., Barnet, C., Olsen, E., Chen, L., & Maddy, E. (2005). On the determination of atmospheric minor gases by the method of vanishing partial derivatives with application to CO₂. *GRL*, *32*, L22803, doi:10.1029/2005GL024165.
- Clough, S., Shephard, M., Mlawer, E., Delamere, J., Iacono, M. J., Cady-Pereira, K., ... Brown, P. (2005). Atmospheric radiative transfer modeling : a summary of the AER codes. *J. Quant. Spectrosc. Rad. Trans.*, *91*, 233–244; doi:10.016/j.qsrt2004.05.058.
- Dee, D., Kallen, E., Simmons, A., & Haimberger, L. (2011). Comments on Reanalyses suitable for characterizing long-term trends. *BAMS*, doi:10.1175/2010BAMS3070.1.
- De Souza-Machado, S., Strow, L. L., Imbiriba, B., McCann, K., Hoff, R., Hannon, S., ... Torres, O. (2010). Infrared retrievals of dust using AIRS: comparisons of optical depths and heights derived for a North African dust storm to

- other collocated EOS A-Train and surface observations. *J. Geophys. Res.*, *115*, doi:10.1029/2009JD012842, D15201.
- De Souza-Machado, S., Strow, L. L., Motteler, H., & Hannon, S. (2020). kCARTA : A Fast Pseudo Line by Line Radiative Transfer Algorithm with Analytic Jacobians, Fluxes, Non-Local Thermodynamic Equilibrium and Scattering. *Atmos. Meas. Tech.*, *31*, 323–339, <https://doi.org/10.5194/amt-13-323-2020>.
- De Souza-Machado, S., Strow, L. L., Tangborn, A., Huang, X., Chen, X., Liu, X., ... Yang, Q. (2018). Single-footprint retrievals for AIRS using a fast TwoSlab cloud-representation model and the SARTA all-sky infrared radiative transfer algorithm. *Atmos. Meas. Tech.*, *11*, 529–550, <https://doi.org/10.5194/amt-11-529-2018>.
- Dessler, A., Zhang, Z., & Yang, P. (2008). Water-vapor climate feedback inferred from climate fluctuations, 2003–2008. *Geophys. Res. Lett.*, *35*, <https://doi.org/10.1029/2008GL035333>.
- Durre, I., Vose, R., & Wuertz, D. (2006). Overview of the Integrated Global Radiosonde Archive. *J. Clim.*, *19*, 53–68; DOI:10.1175/JCLI3594.1.
- Eyring, V., Bony, S., Meehl, G., Senior, C., Stevens, B., Stouffer, A., & Taylor, K. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model. Dev.*, *9*, 1937–1958; doi:10.5194/gmd-9-1937-2016.
- Feng, J., & Huang, Y. (2021). Impacts of tropical cyclones on the thermodynamic conditions in the tropical tropopause layer observed by A-Train satellites. *AMT*, *21*(2-), 15493–15518; <https://doi.org/10.5194/acp-21-15493-2021>.
- Fu, D., Kulawik, S., Bowman, J., K. abd Worden, & Eldering, A. (2018). Retrievals of tropospheric ozone profiles from the synergism of AIRS and OM methodology and validation. *AMT*, *11*, 5587–5605; <https://doi.org/10.5194/amt-11-5587-2018>.
- Gelaro, R., & Coauthors. (2017). MERRA-2 Overview: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2). *J. Clim.*, *30*, 5419–5454, doi: 10.1175/JCLI-D-16-0758.1.
- Gordon, I., & Rothman, L. e. a. (2022). The HITRAN 2020 molecular spectroscopic database. *J. Quant. Spectrosc. Rad. Trans.*, *277*, 1–82; <https://doi.org/10.1016/j.jqsrt.2021.107949>.
- Haiden, T., Dahoui, M., Ingleby, B., de Rosnay, P., Prates, C., Kusc, E., ... Jones, L. (2018). *Use of in situ surface observations at ECMWF* (No. 834).
- Harries, J., Brindley, H., & Geer, A. (1998). Climate variability and trends from operational satellite spectral data. *GRL*, *25*, 3975–3978; <https://doi.org/10.1029/1998GL900056>.
- Held, I., & Soden, B. (2000). Water vapor feedback and global warming. *Annu. Rev. Energy. Environ.*, *25*, 441–475; doi:10.1146/annurev.energy.25.1.441.
- Held, I., & Soden, B. (2006). Robust Responses of the Hydrological Cycle to Global Warming. *J. Clim.*, *19*, 5686–5699; <https://doi.org/10.1175/JCLI3990.1>.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., ... Thépaut, J.-N. (2020). The ERA5 Global Reanalysis. *Quart. J. Roy. Meteorol. Soc.*, *146*, 1999–2049; <https://doi.org/10.1002/qj.3803>.
- Hobbs, W., Klekociuk, A., & Pan, Y. (2020). Validation of reanalysis Southern Ocean atmosphere trends using sea ice data. *ACP*, *20*, 14757–14768; <https://doi.org/10.5194/acp-20-14757-2020>.
- Houghton, J., Jenkins, G. J., & Ephraums, J. J. (1990). *Climate change, the ipcc scientific assessment*. Cambridge University Press.
- Huang, X., Chen, X., Fan, X., Kato, S., Loeb, N., Bosilovich, M., ... Strow, L. (2023). A Synopsis of AIRS Global-Mean Clear-Sky Radiance Trends From 2003 to 2020. *JGR*, *127*, e2022JD037598; <https://doi.org/10.1029/2022JD037598>.
- Huang, Y., Ramaswamy, V., Huang, X., Fu, A., & Bardeen, C. (2007). A strict test in climate modeling with spectrally resolved radiances: GCM simulation versus AIRS observations. *GRL*, *34*, L24707; <https://doi.org/10.1029/2007GL031409>.

- IPCC. (2021). *Climate Change 2021 - The Physical Science Basis*. Cambridge University Press; <https://doi.org/10.1017/9781009157896>.
- Irion, F., Kahn, B., Schreier, M., Fetzer, E., Fishbein, E., Fu, D., . . . Yue, Q. (2018). Single-footprint retrievals of temperature, water vapor and cloud properties from AIRS. *Atmos. Meas. Tech.*, *11*(2), <https://doi.org/10.5194/amt-11-971-2018>.
- Jevanjee, N., Seely, J., Paynter, D., & Fueglistaler, S. (2021). An Analytical Model for Spatially Varying Clear-Sky CO₂ Forcing. *J. Clim.*, *34*, 9463-9480; <https://doi.org/10.1175/JCLI-D-19-0756.1>.
- Kahn, B., Eldering, A., Fetzer, E., Fishbein, E., Lee, S.-Y., Liou, K., . . . Hannon, S. (2005). Nighttime cirrus detection using the Atmospheric Infrared Sounder window channels and total column water vapor. *J. Geophys. Res.*, *110*, doi:0.1029/2004JD005430,D07203.
- Kahn, B., Irion, F., Dang, V., Manning, E., Nasiri, S., Naud, C., . . . Worden, J. (2014). The Atmospheric Infrared Sounder version 6 Cloud Products. *Atmos. Chem. Phys.*, *14*, 399-426.
- Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., . . . Josphe, D. (1996). The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Amer. Met. Soc.*, *77*(D20S93), 437-472; [https://doi.org/10.1175/1520-0477\(1996\)077<0437:TNYRP>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<0437:TNYRP>2.0.CO;2).
- Keeling, C., Bacastow, R., Bainbridge, A., Ekdahl, C., Guenther, P., & Waterman, L. (1976). Atmospheric carbon dioxide variations at mauna loa observatory, hawaii. *Tellus*, *28*, 538-551.
- King, M., Platnick, S., Menzel, P., Ackerman, S., & Hubanks, P. (2013). Spatial and Temporal Distribution of Clouds Observed by MODIS Onboard the Terra and Aqua Satellite. *IEEE*, *51*, 3826-3852, [10.1109/TGRS.2012.2227333](https://doi.org/10.1109/TGRS.2012.2227333).
- Kistler, C., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., . . . Fiorino, M. (2001). The NCEP-NCAR 50-year reanalysis: Monthly means CD-ROM and documentation. *BAMS*, *82*, 247-267; DOI: [10.1175/1520-0477\(2001\)082<0247:tnnyrm>2.3.co;2](https://doi.org/10.1175/1520-0477(2001)082<0247:tnnyrm>2.3.co;2).
- Lambert, A., Read, W., Livesey, N., & Fuller, R. (2021). *MLS/Aura Level 3 Monthly Binned Vapor (H₂O) Mixing Ratio on Assorted Grids V005* (Tech. Rep.). https://disc.gsfc.nasa.gov/datasets/ML3MBH2O_005/summary: NASA Greenbelt, MD, USA, Goddard Earth Sciences Data and Information Services Center (GES DISC).
- Lambert, A., Read, W., Livesey, N., Santee, M., Manney, G., Froidevaux, L., . . . Atlas, B. (2007). Validation of the Aura Microwave Limb Sounder middle atmosphere water vapor and nitrous oxide measurements. *JGR*, *112*, <https://doi.org/10.1029/2007JD008724>.
- LeMarshall, J., Jung, J., Deber, J., Chahine, M., R., T., Lord, S., . . . Tahara, Y. (2006). Improving Global Analysis and Forecasting with AIRS. *Bull. Amer. Met. Soc.*, *87*, 891-894, doi: [10.1175/BAMS-87-7-891](https://doi.org/10.1175/BAMS-87-7-891).
- Lenssen, N., Schmidt, G., Hansen, J., Menne, M., Persin, A., Ruedy, R., & Zyss, D. (2019). Improvements in the GISTEMP uncertainty model. *JGR*, *124*(12), 6307-6326, doi:10.1029/2018JD029522.
- Lin, Y., & Oey, L. (2020). Global Trends of Sea Surface Gravity Wave, Wind, and Coastal Wave Setup. *J. Clim.*, *33*, 769-785, <https://doi.org/10.1175/JCLI-D-19-0347.1>.
- Livesey, N., Van Snder, W., Read, W., & Wagner, P. (2006). Retrieval Algorithms for the EOS Microwave Limb Sounder (MLS). *IEEE*, *44*, 1144-1155.
- Maddy, E., & Barnett, C. (2008). Vertical Resolution Estimates in Version 5 of AIRS Operational Retrievals. *IEEE*, *66*(8), 2375-2384, DOI [10.1109/TGRS.2008.917498](https://doi.org/10.1109/TGRS.2008.917498).
- Manabe, S., & Wetherald, R. T. (1967). Thermal equilibrium of the atmosphere with a given distribution of relative humidity. *J. Atmos. Sci.*, *24*, 241-259.

- Manning, E. (2022). *Nasa jpl private communication, 4/2022*. (AIRS CCR L3 data)
- Manning, E., Aumann, H., Broberg, S., Pagano, T., Wilosn, R., Yanovsky, I., & Strow, L. (2020). *Eos airs l1c data product user guide* (Tech. Rep.). https://docserver.gesdisc.eosdis.nasa.gov/public/project/AIRS/V6.7_L1C_Product_User_Guide.pdf: NASA.
- Masuda, K., Takashima, T., & Takayama, Y. (1988). Emissivity of pure and sea waters for the model sea surface in the infrared window regions. *Remote Sensing of Environment*, 24, 313–329.
- McMillan, W., Barnet, C., Strow, L., Chahine, M., MCCourt, M., Warner, J., . . . Datta, S. (2005). Daily global maps of carbon monoxide from nasa’s atmospheric airs sounder. *Geophys. Res. Lett.*, 32, L11801.
- Mears, C., & Wentz, F. (2009). Construction of the Remote Sensing Systems v3.2 Atmospheric Temperature Records from the MSU and AMSU Microwave Sounders. *J. Atmos. Oceanic Technol.*, 26, 1040–1056, doi: 10.1175/2008JTECHA1176.1.
- Mears, C., & Wentz, F. (2016). Sensitivity of Satellite-Derived Tropospheric Temperature Trends to the Diurnal Cycle Adjustment. *J. Climate*, 29, 3629–3646, <https://doi.org/10.1175/JCLI-D-15-0744.1>.
- Menzel, W., Frey, R., Borbas, E., Baum, B. A., Cureton, G., & Bearson, N. (2016). Reprocessing of HIRS Satellite Measurements from 1980 to 2015: Development toward a Consistent Decadal Cloud Record. *JAMC*, 55, 2397–2410; <https://doi.org/10.1175/JAMC-D-16-0129.1>.
- Muller, R., Kunz, A., Hurst, D., Rold, C., Kramer, M., & Riese, M. (2016). The need for accurate long-term measurements of water vapor in the upper troposphere and lower stratosphere with global coverage. *Earth’s Future*, 4, 25–32; doi:10.1002/2015EF000321.
- Nalli, N., Jung, J., Stegmann, P., Johnson, B., & Zhou, L. (2022). Temperature-dependent optical constants of water in the thermal infrared derived from data archaeology. *Opt. Cont.*, 1, 738–746; <https://doi.org/10.1364/OPTCON.450833>.
- Newman, S., Smith, J., Glew, M., Rogers, S., & Taylor, J. (2005). Temperature and salinity dependence of sea surface emissivity in the thermal infrared. *QJRMSS*, 131, 2539–2557; <https://doi.org/10.1256/qj.04.150>.
- Palchetti, L., Brindley, H., Bantges, R., Buehler, S. A., Camy-Peyret, C., Carli, B., . . . Serio, C. (2020). FORUM: unique far-infrared satellite observations to better understand how Earth radiates energy to space. *Bull.Amer.Met.Soc.*, 101, <https://doi.org/10.1175/BAMS-D-19-0322.1>.
- Peters, W., Jacobson, A., Sweeney, C., & Tans, P. (2007). An atmospheric perspective on North American carbon dioxide exchange: CarbonTracker. *Proc. Nat. Aca. Sci.*, 104, 18925–18930; <https://doi.org/10.1073/pnas.0708986104>.
- Prakash, S., & Norouzi, H. (2020). Land surface temperature variability across India: a remote sensing satellite perspective. *Theor. Appl. Climatol.*, 139, 773–784; <https://doi.org/10.1007/s00704-019-03010-8>.
- Raghuraman, S., Paynter, D., V., R., & Menzel, R. H. X. (2023). Greenhouse Gas Forcing and Climate Feedback Signatures Identified in Hyperspectral Infrared Satellite Observations. *Geophys. Res. Lett.*, 50, <https://doi.org/10.1029/2023GL103947>.
- Rodgers, C. (2000). Inverse Methods for Atmospheric Sounding. In (p. 256 pages). World Scientific, Singapore.
- Roemer, F., Buehler, S., Brath, M., Kluft, L., & John, V. (2023). Direct observation of Earth’s spectral long-wave feedback parameter. *Nature Geoscience*, 16, 416–421; <https://doi.org/10.1038/s41561-023-01175-6>.
- Scherrer, S., de Valk, C., Begert, M., Gubler, S., S., K., & Croci-Maspoli, M. (2024). Estimating trends and the current climate mean in a changing climate. *Clim. Services*, 22, <https://doi.org/10.1016/j.cliser.2023.100428>.

- Schneider, P., T. an O’Gorman, & Levine, X. (2010). Water Vapor and the Dynamics of Changes. *Rev. Geophys.*, *48*, <https://doi.org/10.1029/2009RG000302>.
- Shao, X., Ho, S.-P., Jung, X., Zhou, Z., Chen, Y., Liu, T.-C., . . . Dong, J. (2023). Characterizing the tropospheric water vapor spatial variation and trend using 2007–2018 COSMIC radio occultation and ECMWF reanalysis data. *ACP*, *23*, 14187–14218; <https://doi.org/10.5194/acp-23-14187-2023>.
- Sherwood, S., Ingram, W., Tshushuma, Y., Satoh, M., Roberts, M., Vidale, P., & O’Gorman, P. (2010). Relative humidity changes in a warmer climate. *J. Geophys. Res.*, *115*, <https://doi.org/10.1029/2009JD012585>.
- Shi, L., & Bates, J. (2011). Three decades of intersatellite-calibrated High-Resolution Infrared Radiation Sounder upper tropospheric water vapor. *J. Geophys. Res.*, *116*(D4), D04108, <https://doi.org/10.1029/2010JD014847>.
- Smith, N., & Barnett, C. (2019). Uncertainty Characterization and Propagation in the Community Long-Term Infrared Microwave Combined Atmospheric Product System (CLIMCAPS). *Remote Sensing*, *11*, 1227–1251, doi:10.3390/rs11101227.
- Smith, N., & Barnett, C. (2020). CLIMCAPS observing capability for temperature, moisture, and trace gases from AIRS/AMSU and CrIS/ATMS. *AMT*, *13*, 4437–4459, <https://doi.org/10.5194/amt-13-4437-2020>.
- Smith, N., & Barnett, C. (2023). Practical Implications of CLIMCAPS Cloud Clearing and Derived Quality Metrics. *Earth Space Sci*, *10*, e2023EA002913. <https://doi.org/10.1029/2023EA002913>.
- Sodemann, H. (2019). Beyond Turnover Time: Constraining the Lifetime Distribution of Water Vapor from Simple and Complex Approaches. *JAS*, *77*, 411–433; DOI: 10.1175/JAS-D-18-0336.1.
- Soden, B., & Held, I. (2006). An assessment of climate feedbacks in coupled ocean-atmosphere models. *J. Clim.*, *19*, 3354–3360.
- Strow, L., & DeSouza-Machado, S. (2020). Establishment of AIRS climate-level radiometric stability using radiance anomaly retrievals of minor gases and sea surface temperature. *Atmos. Meas. Tech.*, *13*, <https://doi.org/10.5194/amt-13-4619-2020>.
- Strow, L., Hannon, S., DeSouza-Machado, S., Tobin, D., & Motteler, H. (2003). An overview of the AIRS radiative transfer model. *IEEE Transactions on Geosciences and Remote Sensing*, *41*, 303–313.
- Strow, L., Hannon, S., Weiler, M., Overoye, K., Gaiser, S., & Aumann, H. (2003). Pre-launch spectral calibration of the Atmospheric InfraRed Sounder (AIRS). *IEEE Transactions on Geosciences and Remote Sensing*, *41*, 274–286.
- Strow, L., Hoppelwhite, C., Motteler, H., Buczkowski, S., & DeSouza-Machado, S. (2021). A Climate Hyperspectral Infrared Radiance Product (CHIRP) Combining the AIRS and CrIS Satellite Sounding Record. *Rem. Sens.*, *13*(3), <https://doi.org/10.3390/rs13030418>.
- Susskind, J. (2006). Contribution of the AIRS Shortwave Sounding Channels to Retrieval Accuracy. In *Algorithms and technologies for multispectral, hyperspectral, and ultraspectral imagery xii*, vol. 6233, orlando, fl.
- Susskind, J., Barnett, C., & Blaisdell, J. (2003). Retrieval of atmospheric and surface parameters from AIRS/AMSU/HSB data under cloudy conditions. *IEEE Transactions on Geosciences and Remote Sensing*, *41*, 390–409.
- Susskind, J., Blaisdell, J., & Iredell, L. (2014). Improved methodology for surface and atmospheric soundings, error estimates, and quality control procedures: the Atmospheric Infrared Sounder science team version-6 retrieval algorithm. *J. App. Rem. Sens.*, *8*(1), 084994 [10.1117/1.JRS.8.084994].
- Teixeira, J., Wilson, R., & Thrastarson, H. (2024). Direct observational evidence from space of the effect of CO₂ increase on longwave spectral radiances: the unique role of high-spectral-resolution measurements. *ACP*, *24*, 6375–6383; <https://doi.org/10.5194/acp-24-6375-2024>.

- Thorne, P., Lanzante, J., Peterson, T., Seidel, D. J., & Shine, K. (2011). Tropospheric Temperature Trends : history of an ongoing controversy. *Wiley Interdisciplinary Reviews : Climate Change*, 2, 66-88.
- Tian, B., E., M., Roman, J., Thrastarson, H., & Fetzer, R., E. ad Monarrexz. (2020). *Airs version 7 level 3 product user guide* (Tech. Rep.). Jet Propulsion Laboratory, California Institute of Technology. Retrieved from https://docserver.gesdisc.eosdis.nasa.gov/public/project/AIRS/V7_L3_User_Guide.pdf
- van der Ent, R. J., & Tuinenburg, O. A. (2017). The residence time of water in the atmosphere revisited. *Hydrol. Earth Syst. Sci.*, 21, 779–790, <https://doi.org/10.5194/hess-21-779-2017>.
- Warner, J., Wei, Z., Strow, L., Dickerson, R. R., & Nowak, J. (2016). The global tropospheric ammonia distribution as seen in the 13-year AIRS measurement record. *AMT*, 16, 5467–5479;doi:10.5194/acp-16-5467-2016.
- Whitburn, S., Clarisse, L., Bouillon, M., Safieddine, S., George, M., Dewitte, S., ... Clerbaux, C. (2021). Trends in spectrally resolved outgoing longwave radiation from 10 years of satellite measurements. *npj Clim Atmos Sci*, 4, 48, <https://doi.org/10.1038/s41612-021-00205-7>.
- Wong, A., Wijffels, S., Riser, S., et al. (2020). Argo Data 1999–2019: Two Million Temperature-Salinity Profiles and Subsurface Velocity Observations From a Global Array of Profiling Floats. *Front. Mar. Sci.*, 7, <https://doi.org/10.3389/fmars.2020.00700>.
- Wu, W., Liu, X., Lei, X., L. amd Xiong, Yang, Q., Yue, Q., Zhou, D., & Larar, A. (2023). Single field-of-view sounder atmospheric product retrieval algorithm : establishing radiometric consistency for hyper-spectral sounder retrievals. *AMT*, 16, 4807–4832, <https://doi.org/10.5194/amt-16-4807-2023>.
- Zhou, D., Larar, A., Liu, X., Smith, W., Strow, L., Yang, P., ... Calbet, X. (2011). Global Land Surface Emissivity Retrieved From Satellite Ultraspectral IR Measurements. *IEEE Trans. Geosci. Remote Sens.*, 49(4), 1277-1290.
- Zou, M., Xiong, X., Wu, Z., Li, S., Zhang, Y., & Chen, L. (2019). Increase of Atmospheric Methane Observed from Space-Borne and Ground-Based Measurement. *Rem. Sens.*, 11(8), 16; <https://doi.org/10.3390/rs11080964>.